

AHA SCIENTIFIC STATEMENT

Cardiovascular Health in African Americans

A Scientific Statement From the American Heart Association

Endorsed by the American College of Cardiology

BACKGROUND AND PURPOSE: Population-wide reductions in cardiovascular disease incidence and mortality have not been shared equally by African Americans. The burden of cardiovascular disease in the African American community remains high and is a primary cause of disparities in life expectancy between African Americans and whites. The objectives of the present scientific statement are to describe cardiovascular health in African Americans and to highlight unique considerations for disease prevention and management.

METHOD: The primary sources of information were identified with PubMed/Medline and online sources from the Centers for Disease Control and Prevention.

RESULTS: The higher prevalence of traditional cardiovascular risk factors (eg, hypertension, diabetes mellitus, obesity, and atherosclerotic cardiovascular risk) underlies the relatively earlier age of onset of cardiovascular diseases among African Americans. Hypertension in particular is highly prevalent among African Americans and contributes directly to the notable disparities in stroke, heart failure, and peripheral artery disease among African Americans. Despite the availability of effective pharmacotherapies and indications for some tailored pharmacotherapies for African Americans (eg, heart failure medications), disease management is less effective among African Americans, yielding higher mortality. Explanations for these persistent disparities in cardiovascular disease are multifactorial and span from the individual level to the social environment.

CONCLUSIONS: The strategies needed to promote equity in the cardiovascular health of African Americans require input from a broad set of stakeholders, including clinicians and researchers from across multiple disciplines.

Mercedes R. Carnethon, PhD, FAHA, Chair
Jia Pu, PhD
George Howard, DrPH, FAHA
Michelle A. Albert, MD, MPH, FAHA
Cheryl A.M. Anderson, PhD, FAHA
Alain G. Bertoni, MD, MPH, FAHA
Mahasin S. Mujahid, PhD
Latha Palaniappan, MD, MS, FAHA
Herman A. Taylor Jr, MD, FAHA
Monte Willis, MD, PhD, FAHA
Clyde W. Yancy, MD, FAHA
On behalf of the American Heart Association Council on Epidemiology and Prevention; Council on Cardiovascular Disease in the Young; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Functional Genomics and Translational Biology; and Stroke Council

Key Words: AHA Scientific Statements ■ African Americans ■ cardiovascular diseases ■ disease management ■ prevention and control ■ risk factors

© 2017 American Heart Association, Inc.

Despite advances in the identification of risk factors for cardiovascular disease (CVD) and the widespread use of evidence-based strategies to manage CVD, racial/ethnic disparities in CVD morbidity and mortality persist in the United States. Across nearly every metric, African Americans have poorer overall cardiovascular health than non-Hispanic whites, and CVD mortality is higher in African Americans than whites.^{1,2} In fact, little has changed since 2005 when notable disparities in prevalence, disease management, and outcomes were reported in a special issue of *Circulation*.³ The American Heart Association (AHA) is a leader in highlighting disparities in CVD by race and ethnicity. The present scientific statement on cardiovascular health in African Americans follows statements published on Asian Americans⁴ and US Hispanic/Latinos.⁵ Our intention is for the statement to be used by clinicians, public health practitioners, and policy makers to interrupt these adverse trends and to move toward cardiovascular health equity for African Americans.

SIGNIFICANCE AND RATIONALE

African Americans are the oldest nonnative racial groups in the United States, with the large initial influx coming to America involuntarily during the transatlantic slave trade. Since that time, individuals of African descent from around the world (eg, Africa, the Caribbean, Latin America) have immigrated to the United States and contribute to the diversity of language, customs, and cultures of the US African American population. In the present report, we have chosen the term African American to refer to black Americans of African descent living in the United States. However, when referencing work from studies that specifically denote the inclusion of non-Hispanic blacks, we have retained their original label. At present, African Americans make up 13.3% of the US population and are the second largest racial/ethnic minority behind Hispanic/Latinos.⁶

In 2012, the life expectancy of African Americans was 3.4 years shorter than that of whites (75.5 versus 78.9 years, respectively). The contrasts are most striking when studied by race and sex: White women have the longest life expectancy at 81.4 years, followed by black women at 78.4 years, white men at 76.7 years, and black men at 72.3 years.⁷ Among the 25 leading causes of death, 6 of the 10 diseases that are substantial contributors to years of life lost are CVD risk factors (ie, hypertension, diabetes mellitus, renal disease) or CVDs (ie, ischemic heart disease, heart failure, and stroke). In the most recent report by the Centers for Disease Control and Prevention, CVDs were estimated to explain 32% of the mortality difference between African American and white men and 43% of the difference between African American and white women in 2009.⁸ Together, these conditions contributed to >2.0 million years of

life lost in the African American population between 1999 and 2010.

OBJECTIVES

The objectives of the current statement are to describe cardiovascular health and the burden of CVD in the population; to discuss the contribution of traditional CVD risk factors and adverse health behaviors to disparities in cardiovascular health between African Americans and whites; to describe the contribution of comorbidities that are overrepresented among African Americans to CVD; to identify and discuss genetic and biological mechanisms that might contribute to the disease pathways leading to CVD in African Americans; to highlight unique considerations (ie, differential effects of pharmacological strategies) in disease prevention and management in African Americans; and to discuss the social, cultural, and environmental factors that influence prevention and disease management in African Americans. Aside from a brief discussion about the origin of disparities in youth, the current statement focuses on health in adults. A detailed discussion of strategies to reduce disparities in CVDs between African Americans and other racial/ethnic groups is beyond the scope of the present statement. We conclude the statement by recommending a broad set of strategies for change, from additional research to workforce development.

BURDEN OF CVDs AND STROKE

The term *cardiovascular diseases* will collectively include coronary heart disease (CHD), sudden cardiac death/sudden cardiac arrest, stroke/transient ischemic attack, and peripheral arterial disease. The AHA has played a key role in summarizing trends in cardiovascular health and disease in the annual AHA statistical report.² Because these data are published and updated annually, we only briefly summarize the burden of these diseases in African Americans compared with non-Hispanic whites.

Coronary Heart Disease

Although the rates of CHD have declined in recent decades, those declines are smaller among African Americans than whites. In the ARIC study (Atherosclerosis Risk in Communities), the decline in CHD incidence among African American men was half (−3.2%/y) that of the decline among white men (−6.5%/y). African American women experienced a decline of −4.0%/y, whereas white women experienced a decline of −5.2%/y.¹ In 2010 (the most recent year that national prevalence rates were available), the self-reported prevalence of

diagnosed CHD was 6.5% in African Americans compared with 5.8% in whites, a difference that was not statistically significant. However, the modestly higher rate in African Americans is driven by the excess among African American women (5.9%) compared with white women (4.0%); among men, rates were higher among whites (7.7%) versus African Americans (7.3%).⁹ Longitudinal data from the REGARDS study (Reasons for Geographic and Racial Differences in Stroke) describe no difference in incident CHD between African American and white men (hazard ratio [HR], 1.04; 95% confidence interval [CI], 0.84–1.29) and only a marginally and nonsignificantly higher incidence in African American women versus white women (HR, 1.25; 95% CI, 0.96–1.62). However, African American men and women have substantially higher rates of fatal CHD than whites (men: HR, 2.18; 95% CI, 1.24–2.56; women: HR, 1.63; 95% CI, 1.02–2.62).¹⁰

Heart Failure

The incidence, prevalence, and prognosis of heart failure are less favorable among African Americans and are largely attributable to the higher burden of traditional risk factors among African Americans. Disparities in the incidence of heart failure are most prominent at young ages, as reported by the CARDIA study (Coronary Artery Risk Development in Young Adults), in which 26 of the 27 incident heart failure cases that occurred in individuals <50 years of age were among black participants.¹¹ Although disparities in heart failure persist in middle-aged and older adults (mean age, 62 years) in MESA (Multi-Ethnic Study of Atherosclerosis; HR, 1.81; 95% CI, 1.07–3.07 in African Americans versus whites), statistical adjustment for established risk factors (ie, age, sex, diabetes mellitus, hypertension, cholesterol, smoking status, and left ventricular hypertrophy) explained all of the excess risk among African Americans versus whites (HR, 1.42; 95% CI, 0.81–2.48).¹² Findings were similar in the ARIC study, but the extended follow-up (15 versus 4 years in MESA) and large number of total events (n=1282) provided additional evidence that the disparities were present with younger age at disease onset. Among men, adjustment for established risk factors eliminated any differences in heart failure incidence between African Americans and whites (HR, 0.86; 95% CI, 0.70–1.06, African Americans versus whites). However, among women, African Americans were significantly more likely to experience events within the first 7.5 years even after statistical adjustment for risk factors (HR, 1.79; 95% CI, 1.25–2.55 versus white women). The disparity between African American and white women was attenuated only after adjustment during the second half of follow-up when women were older (HR, 0.93; 95% CI, 0.46–1.90, African American versus white). In that same study, the age-adjusted 30-day case

fatality rate (per 1000 person-years) was significantly higher ($P<0.05$) in African American men (51.8; 95% CI, 44.1–59.4) and women (46.1; 95% CI, 39.8–52.5) compared with white men (41.2; 95% CI, 36.9–45.6) and white women (35.8; 95% CI, 30.6–41.4).¹³

Sudden Cardiac Arrest/ Sudden Cardiac Death

Sudden cardiac arrest, a sudden pulseless condition frequently attributable to underlying cardiac causes, has high fatality rates outside of hospital settings (ie, sudden cardiac death). Both sudden cardiac arrest and sudden cardiac death are higher in African Americans compared with whites, primarily because of a higher burden of traditional and nontraditional (eg, sickle cell trait) CVD risk factors in African Americans.¹⁴ In the Oregon Sudden Unexpected Death Study, a community-based epidemiological study initiated in 2002 to collect information about out-of-hospital cardiac arrest, African Americans were twice as likely to experience sudden cardiac death. From 2002 to 2012, the rate of sudden cardiac arrest was 175 per 100 000 in African American men compared with 84 per 100 000 in white men. African American women experienced sudden cardiac death at a rate of 90 per 100 000, whereas white women experienced sudden cardiac arrest at a rate of 40 per 100 000. African American men and women who experienced sudden cardiac arrest were on average >6 years younger than their white counterparts. According to the National Registry on Cardiopulmonary Resuscitation, when patients are hospitalized for sudden cardiac arrest, African Americans are less likely to survive to discharge (25.2%) than whites (37.4%).¹⁵

Cerebrovascular Disease/Stroke

Cerebrovascular disease incidence and mortality, inclusive of transient ischemic attacks, ischemic stroke, and intracerebral hemorrhage, are notably higher in African Americans compared with whites in the United States. Notably, although stroke mortality has fallen by 80% across all ages over the past 60 years, there has been no meaningful decrease in the magnitude of the African American to white racial disparity in stroke mortality.¹⁶ Since the earliest studies in the 1950s, stroke mortality rates in nonwhites (predominately African Americans) remain 4.5-fold higher than among whites. Whether the higher stroke mortality in African Americans is attributable to a higher incidence of stroke, higher case fatality after stroke events, or a combination is not entirely known. In the ARIC study, the incidence rate ratio for stroke comparing African Americans to whites <55 years of age was modestly larger (relative risk [RR], 2.77; 95% CI, 1.37–5.62) than the same comparison in those >55 years of age (RR, 2.23; 95% CI, 1.66–3.00).¹⁷

However, most African Americans in ARIC were recruited from a single clinical site in the southern Stroke Belt region (Jackson, MS), confounding the racial and geographic disparities in stroke risk and complicating the interpretation of these results. The implication of the regional differences in the racial disparity in magnitude is that the Stroke Belt is more “potent” for African Americans than whites, and as a corollary, it is possible that between 2% and 13% of the black-white difference in stroke risk (and likely coronary risk) is a result of confounding with geographic differences in risk.¹⁸

Data from both REGARDS and GCKSS (Greater Cincinnati/Northern Kentucky Stroke Study) have shown that age-related differences in the magnitude of racial disparity in stroke mortality are mirrored by age-related changes in the magnitude of the racial disparity in stroke incidence. GCKSS reported patterns of stroke incidence in 1999 in the Cincinnati region that were similar to the national pattern for stroke mortality, with stroke mortality 2.6 times greater in African Americans than whites at 45 to 54 years of age but decreasing to 1.8 times for 55 to 64 years of age, 1.2 times for 65 to 74 years of age, 0.9 times for 75 to 84 years of age, and 0.8 times for ≥ 85 years of age.¹⁹ In the same report, stroke case fatality was 24% lower in African Americans than whites, an observation that was relatively consistent across stroke subtypes of infarction and hemorrhage.¹⁹

There is a strong age-related difference in the risk of intracerebral hemorrhage in African Americans compared with whites. At 55 to 74 years of age, African Americans were 1.8 times more likely to experience intracerebral hemorrhage, but that difference was only modestly greater (RR, 1.23 times) for ages ≥ 75 years.²⁰ A pooled analysis of ARIC and the CHS (Cardiovascular Health Study) showed a similar pattern whereby the RR of intracerebral hemorrhage for African Americans versus whites was 5.8 at 45 years, 1.7 at 65 years, and 0.94 at 75 years of age.²¹ Among 45- to 64-year-olds in REGARDS, the incidence rate of intracerebral hemorrhage (per 100 000) was doubled for African Americans (46.0; 95% CI, 26.5–79.7) compared with whites (21.5; 95% CI, 11.2–41.2). However, these associations were reversed in older adults. The risk of incident intracerebral hemorrhage was 40.1 (95% CI, 17.8–90.4) in African Americans and 65.1 (95% CI, 39.1–108.4) in whites 65 to 74 years old. At ages ≥ 75 years, the incidence rates were 65.8 (95% CI, 24.4–177.8) in African Americans and 105.0 (95% CI, 64.3–171.3) in whites.²²

Peripheral Arterial Disease

Atherosclerotic disease affecting the arteries and vessels outside of the heart, peripheral arterial disease/peripheral vascular disease, is a common geriatric disease with a prevalence of 12% to 20% among adults >80 years

of age.²³ Across the age range, the rate of peripheral arterial disease is twice as high in African Americans compared with whites.²⁴ Traditional risk factors, namely cigarette smoking, diabetes mellitus, and hypertension, are the strongest risk factors for peripheral arterial disease, but statistical adjustment for these and other traditional risk factors does not completely eliminate the excess prevalence in African Americans compared with whites.^{25,26} In the MESA study, the adjusted odds for incident peripheral arterial disease were 1.67 times higher in African Americans compared with whites.²⁷ The San Diego Population Study included markers of inflammation in multivariable models to test whether they explained the residual excess risk in African Americans compared with whites but found that, although the RRs for peripheral artery disease between African Americans and whites were further attenuated, they remained statistically significantly higher, ranging from 1.5 to 2.0.²⁸

Although peripheral arterial disease is not considered a direct cause of mortality but rather a reflection of the overall burden of CVD, peripheral arterial disease was listed as the underlying cause of death for 59 681 deaths in 2014. The age-adjusted death rate resulting from peripheral arterial disease was higher among African American men (24.8 per 100 000) than white (19.9), American Indian or Alaska Native (20.8), Hispanic (15.4), or Asian Pacific Islander (8.5) men. Similar patterns of peripheral arterial disease mortality were observed for African American women (16.5) compared with white (13.8), American Indian or Alaska Native (16.1), Hispanic (10.7), and Asian or Pacific Islander (6.8) women.²

Summary

There are marked disparities in the onset of heart failure, stroke, and peripheral vascular disease between African Americans and whites, whereas rates of CHD are not significantly different, particularly among men. However, mortality from all CVDs is significantly higher in African Americans compared with whites, which suggests a role for health care to mitigate disparities with comprehensive screening, an enhanced specificity of diagnoses, and tailored disease management. The prominence of disparities in the onset of CVD at younger ages highlights the contribution of cardiovascular risk factors and adverse health behaviors among African Americans.

TRADITIONAL CVD RISK FACTORS

The AHA 2020 Strategic Impact Goals for Cardiovascular Health Promotion and Disease Reduction provided metrics to determine adherence to current recommendations for CVD prevention. Subsequent reports

identified African American children, adolescents, and adults^{30,31} as less likely than other racial/ethnic groups to achieve ideal cardiovascular health. The following sections describe the burden of cardiovascular risk factors, proposed risk factor management, and adverse health behaviors.

Cardiovascular Risk Factors

Hypertension

Hypertension is arguably the most potent risk to the cardiovascular health of African Americans, as well as the greatest area of opportunity for the prevention of disease if effectively managed and prevented. The prevalence of diagnosed and undiagnosed hypertension among African American men (42.4%) and women (44%) ≥ 20 years of age in the United States³² is among the highest in the world where the population prevalence of hypertension is the highest in low- to middle-income countries (29%–31%).³³ An analysis of trends indicates that rates of hypertension among African Americans remained $\approx 10\%$ to 12% higher than rates among non-Hispanic whites and Mexican Americans since 1999 to 2000 (the year that the National Health and Nutrition Examination Survey [NHANES] became semiannual).³² Percent of African admixture in African Americans and other racial/ethnic groups is positively associated with blood pressure (BP) levels and the prevalence of hypertension.^{34,35}

The origins of adult differences in hypertension begin in youth. African American boys and girls have higher BP levels and a higher prevalence of hypertension (13.8% in African Americans versus 8.4% in whites and 10.4% in Hispanics).³⁶ Findings from the Bogalusa Heart Study indicate that higher BP levels during childhood track into elevated BP in adults.³⁷ These differences persist into older ages, as evidenced by the MESA study, in which the odds of hypertension were 1.5 times higher in African Americans than in whites through age 75 years.³⁸ In REGARDS, the RR of incident hypertension was 1.24 (95% CI, 1.12–1.37) times higher in African American men compared with white men across the life span. In contrast, an interaction with age was observed for women ($P=0.08$). The RR for incident hypertension was significantly higher for African American women 65 to 74 years of age (RR, 1.44; 95% CI, 1.24–1.66) but was not significant for age >75 years (RR, 1.18; 95% CI, 0.84–1.65, African American versus white women).³⁹

On a positive note, African Americans were more likely than whites or Hispanics to be aware of their hypertension and to have it treated.^{40,41} In NHANES, 87% of African Americans, 81% of whites, and 77% of Hispanics were aware of their hypertension, and 80% of African Americans were treated with medications compared with 77% of whites and 70% of Hispanics.⁴⁰

African Americans in REGARDS were also more likely to be aware of their hypertension (odds ratio [OR], 1.45; 95% CI, 1.24–1.71 versus whites) and treated (OR, 1.56; 95% CI, 1.34–1.83, African Americans versus whites).⁴² Despite these favorable trends in awareness and treatment as noted in NHANES, fewer African Americans achieve BP control (47.9%) than non-Hispanic whites (56%).⁴⁰ Similarly, the adjusted odds of hypertension control in REGARDS are lower in African Americans compared with whites (OR, 0.67; 95% CI, 0.60–0.74).⁴²

The prevalence of hypertension in African Americans has significant implications for mortality. The magnitude of the association between systolic BP (SBP) levels and stroke risk is 3 times greater in African Americans than in whites; a 10-mmHg difference in SBP is associated with an 8% (95% CI, 0–16) increase in the stroke risk in whites but a 24% (95% CI, 14–35) increase in African Americans. Within strata of SBP, stroke risk goes up with an increasing number of classes of antihypertensive medications used to treat high BP (RR, 1.42 for 1 class up to 2.48 for ≥ 3 classes).⁴³ Even when treatment recommendations are followed for African Americans, stroke risks remain elevated, suggesting primordial prevention as the best strategy to eliminate the risks of hypertension-related vascular outcomes.

The National Heart, Lung, and Blood Institute (NHLBI) Working Group on Research Needs to Improve Hypertension Treatment and Control in Africans produced a brief report summarizing some of the above findings and, above all, advocating for additional research into the sources of disparities in hypertension control in African Americans, given its contribution to significant disparities in CVDs.⁴⁴ Consequently, addressing disparities in hypertension incidence and management is a high priority.

Diabetes Mellitus

More than 95% of the cases of diabetes mellitus are classified as type 2,⁴⁵ and the combined prevalence of diagnosed and undiagnosed type 2 diabetes mellitus is 14.3% overall but 21.8% in African Americans and 11.3% in non-Hispanic whites according to NHANES 2011 to 2012.⁴⁶ More than 1 in 3 (37%) African Americans with diabetes mellitus were not diagnosed.⁴⁶ The prevalence of diabetes mellitus among African Americans has increased dramatically in the past decades, from 8% in 1988 to 1994 to the current rates.⁴⁶ Similarly, prediabetes increased among African Americans from 14% in 1988 to 1994 to 20% in 2005 to 2010.⁴⁷ The elevated incidence of diabetes mellitus among African Americans extends over the life course, with no age-related decline in the disparity. Over a lifetime, African American men develop diabetes mellitus 1.52 times (95% CI, 1.31–1.78) more often than white men, and African American women are 2.14 (95% CI, 1.86–2.46)

times more likely to develop diabetes mellitus than white women.³⁹

Although type 1 diabetes mellitus remains the most common type of diabetes mellitus among children and adolescents (age <21 years), rates of type 2 diabetes mellitus among youth have increased secondary to the obesity epidemic. African American adolescents (age, 12–19 years) are significantly more likely to develop diabetes mellitus than whites, and their rate falls short of only the rate among American Indian adolescents.⁴⁸ These patterns are particularly troublesome given the long lifetime of exposure to higher glucose levels among African Americans that contribute to mortality and vascular complications of diabetes mellitus.

The diagnosis of diabetes mellitus is made by a combination of fasting glucose, postchallenge glucose, and hemoglobin A_{1c}.⁴⁵ However, relying solely on hemoglobin A_{1c} could underestimate the prevalence of diabetes mellitus in African Americans because of disorders such as sickle cell trait that occur among individuals of African ancestry. Recent research indicates that at a given level of fasting glucose, hemoglobin A_{1c} is statistically significantly lower (5.72%) among those with sickle cell trait versus those without (6.01%).⁴⁹ Delays in diagnosing diabetes mellitus have adverse implications for the development of vascular complications.

African Americans are less likely to be aware of their diabetes mellitus and, when treated, are less likely to achieve adequate control according to common quality metrics defined by the Accountable Care Organization (hemoglobin A_{1c} <9%).⁵⁰ Only 54% of African Americans achieved targets compared with 61% of whites,⁴⁷ which may contribute directly to the elevated excess in microvascular complications of diabetes mellitus in African Americans compared with whites.⁵¹ The age-adjusted death rate among people with diabetes mellitus in 2011 was 40 per 100 000 in African Americans compared with 19 per 100 000 in whites and 26 per 100 000 in Hispanics.⁵² Compared with their white peers, African Americans with diabetes mellitus are 4 times more likely to have visual impairment (caused by diabetic retinopathy)⁵³ and 3.8 times more likely to have end-stage renal disease (resulting from diabetic nephropathy)^{54,55} but possibly less likely to experience lower extremity amputation.⁵⁶

Lipid Disorders

Despite higher atherosclerotic CVD (ASCVD) rates and higher mortality from CHD among African Americans, lipid profiles among African Americans according to national prevalence estimates are comparable to or lower than those of non-Hispanic whites. For example, the prevalence of elevated total cholesterol (≥ 200 mg/dL) was 37.0% in non-Hispanic white males versus 32.6% in non-Hispanic blacks in NHANES 2011 to 2014. Comparable percentages of elevated total cholesterol among

women were 43.4% among non-Hispanic whites and 36.1% in non-Hispanic blacks.² High-density lipoprotein (HDL) cholesterol is known to be higher in African Americans, and in recent estimates from NHANES, those patterns hold, with the prevalence of low HDL (<40 mg/dL) at 28.4% in non-Hispanic white men compared with only 20.7% in non-Hispanic black men. There were small differences in the prevalence among women (10.3% in non-Hispanic white women and 8% in non-Hispanic black women).² The prevalence of dyslipidemia was higher among African Americans in the JHS (Jackson Heart Study), a cohort in Jackson, MS, compared with national estimates. One third of participants 35 to 84 years of age had hypercholesterolemia, defined as total cholesterol ≥ 240 mg/dL, low-density lipoprotein (LDL) cholesterol (LDL-C) ≥ 160 mg/dL, or triglycerides ≥ 200 mg/dL.⁵⁷ In this population, the most common dysfunction was in elevated LDL-C (18.3%), followed by total cholesterol (15.2%) and then triglycerides (5.4%).⁵⁷ Relying on the lipid panel and a focus on dyslipidemia may underestimate CVD risk in African Americans given the relatively lower likelihood of high LDL-C and triglycerides in African Americans. However, incidence data on dyslipidemia present a different picture.

Over the age of 45, there is a higher incidence of dyslipidemia (total cholesterol ≥ 240 mg/dL, LDL ≥ 160 mg/dL, HDL ≤ 40 mg/dL, or use of lipid-lowering medications) in African American men (RR, 1.15; 95% CI, 1.04–1.28) and women (RR, 1.17; 95% CI, 1.08–1.28) than in their white counterparts. However, among both men ($P=0.10$) and women ($P=0.02$), the disparities become even more pronounced in older ages, with the RR of dyslipidemia in African American men compared with white men increasing from 1.15 (95% CI, 0.82–1.61) at 45 to 54 years of age to 1.26 (95% CI, 1.06–1.51) at 65 to 74 years of age. The RR of dyslipidemia for African American women versus white women is 0.97 (95% CI, 0.77–1.22) at 45 to 54 years of age but 1.39 (95% CI, 1.00–1.95) among women ≥ 75 years of age.³⁹ The apparent contradiction in findings between prevalence and incidence may be explained by the higher rates of CVD mortality among African Americans compared with whites. Disease prevalence is determined on the basis of a combination of disease incidence and the average duration of disease. If African Americans are more likely to die of CVD that could be attributed to dyslipidemia, then the burden of dyslipidemia would not be captured in prevalence estimates because they are exiting the denominator before their disease is captured. Consequently, dyslipidemia management is critically important.

Although the introduction of statin therapy has revolutionized the treatment of lipid disorders, the introduction of therapy requires CVD risk assessment. At the baseline examination of the JHS in 2000 to 2003,

69.7% of participants were aware that they had hypercholesterolemia, 43% of those were being treated, and among those, 88% were controlled.⁵⁷ In the REGARDS study, African Americans were less likely to be aware of their dyslipidemia (OR, 0.69; 95% CI, 0.61–0.78), less likely to have their dyslipidemia treated (OR, 0.77; 95% CI, 0.67–0.89), and if treated, less likely to have their lipids under control (OR, 0.67; 95% CI, 0.58–0.77).⁵⁸

Findings from MESA suggested that guideline-recommended treatment was underused in all ethnic groups examined.⁵⁹ The lowest control rates of dyslipidemia among those on lipid-lowering therapy were seen among African American women (65.7% controlled versus 86.2% for non-Hispanic white women), followed by African American and Hispanic men (both 68.5% versus 76.4% in non-Hispanic white men).⁵⁹ Although the guidelines have been updated to eliminate numerical targets as therapeutic goals,⁶⁰ the historical pattern of undertreatment among African Americans^{57,59} raises the concern that without specific attention by providers and adherence by patients, CVD risk tied to dyslipidemia will remain higher among African Americans than other Americans.

Obesity

Across the age spectrum, obesity rates are higher among African Americans than whites. One in 5 (20%) African American children 2 to 19 years old were obese (defined by a body mass index [BMI] for age value \geq 95th percentile of the 2000 Centers for Disease Control and Prevention growth charts) compared with 15% of whites. The rates of extreme obesity (BMI for age value \geq 120th percentile of the 2000 Centers for Disease Control and Prevention growth charts) in children were more than double in African American children (9%) compared with whites (4%).^{61,62} Among adults \geq 20 years of age, African American women had the highest rates of obesity (BMI >30 kg/m²) at 58%, followed by African American men (38%), white men (34%), and white women (33%).² The prevalence of severe obesity (BMI ≥ 40 kg/m²) among African Americans (12.1%) was double that of the next highest groups (Hispanics, 5.8%, and whites, 5.6%).⁶²

The obesity paradox, the observation of a higher risk of mortality in leaner and normal-weight individuals than among adults who are overweight or have class I obesity,⁶¹ may warrant investigation among African Americans. Previous findings from a large cohort study, the Cancer Prevention Study II, showed only a moderately elevated risk of all-cause mortality with increased weight among African Americans but a much stronger finding among whites.⁶³ There are several potential explanations for the observation of an obesity paradox that fall outside of the scope of the present review (eg, selection bias, reverse causation, residual confounding, and measurement error⁶⁴). One that is potentially most

relevant to African Americans is error arising from the use of BMI to represent adiposity.

Although the most common metric to define obesity clinically and at the population level is BMI, body composition and body fat distribution are more precise indicators of metabolic and cardiovascular risk. NHANES captured waist circumference and found that African American women have a larger waist circumference compared with white women but that there were no differences between African American and white men.⁶⁵ Smaller studies that have direct measures of adiposity via imaging (ie, dual x-ray absorptiometry or computed tomography) describe contradictory findings for body composition and adiposity distribution. In studies that have images of adiposity distribution within regions of the body, African American men and women have less metabolically active abdominal visceral adipose tissue compared with whites after adjustment for total body fat.^{66–68} Recent findings from the Pennington Center Longitudinal Study confirm prior observations and report that African American women and men had significantly higher subcutaneous adipose tissue, which is considered protective, compared with white women and men.^{69,70} However, another study found that these racial differences were reversed after adjustment for total body fat, indicating that African American men had more subcutaneous adipose tissue at a given level of total body fat.⁷⁰

Qualitative and quantitative research on the ideal body size and shape in African Americans describes cultural attitudes that favor a larger body size, particularly for women.⁷¹ These attitudes among African Americans complicate the acknowledgement of awareness about obesity and willingness to engage in weight management programs.^{72,73} In the CARDIA study, obese women who perceived themselves as obese lost 0.09 BMI units annually over 13 years compared with obese women who perceived themselves as normal weight, who gained 0.31 BMI units annually ($P<0.001$).⁷⁴ In a weight-loss study, African American women made a similar number of attempts to lose weight but set a weight-loss goal that was 10 lb higher than the goal set by equally obese white women.⁷⁵

Atrial Fibrillation

Although atrial fibrillation has long been recognized as a potent risk factor for stroke,^{76,77} data from REGARDS recently documented that atrial fibrillation is also a potent risk factor for myocardial events,⁷⁸ a finding confirmed in ARIC⁷⁹ and in a meta-analysis of observational studies and clinical trials.⁸⁰ Despite African Americans having more risk factors for the development of atrial fibrillation, studies have consistently documented a lower prevalence of either self-reported or electrocardiographically defined atrial fibrillation, the so-called atrial fibrillation paradox.⁸¹ The incidence of

atrial fibrillation increases dramatically with age, but the risk of incident atrial fibrillation is ≈ 0.20 to 0.50 times lower in African Americans than in their white counterparts across the adult age spectrum.^{39,82,83} Data from REGARDS suggest that these benefits of a lower prevalence (and incidence) of atrial fibrillation among African Americans are offset by a much lower odds of awareness of atrial fibrillation (OR, 0.32; 95% CI, 0.20–0.52) and the fact that, if aware, African Americans are much less likely to be on treatment with warfarin (OR, 0.28; 95% CI, 0.13–0.60).⁸⁴

Adverse Health Behaviors

Poor Diet Quality

The current guidelines for lifestyle management from the AHA and American College of Cardiology to reduce cardiovascular risk include consuming a dietary pattern that emphasizes fruits, vegetables, and whole grains; includes low-fat dairy products, poultry, fish, legumes, nontropical vegetable oils, and nuts; and limits intake of sweets, sugar-sweetened beverages, and red meats.⁶⁰ It is difficult for all Americans to meet these dietary recommendations, and adherence has been documented as low.⁸⁵ African Americans face unique challenges to adherence to recommendations. One such challenge is food intake preferences that align with a cultural tradition of “soul food.” The traditional soul food diet has components that are healthful and many components that are suboptimal.⁸⁶ An example of a healthful component is the inclusion of many fruits and vegetables such as collard greens, sweet potatoes, tomatoes, dried beans and peas, watermelon, blackberries, corn, and okra. Conversely, the diet can also be described as being high in added fats, sugars, and sodium, with prominent use of high-fat meats for main dishes and the use of deep frying and other cooking techniques that add excess calories and sodium.

Regardless of their geographic residence in the United States, REGARDS has shown that African Americans are much more likely to consume a “southern diet.”⁸⁷ When adherence to a southern diet was categorized into quartiles, only 9% of African Americans fell within the lowest quartile of adherence, and 60% fell within the uppermost quartile. The HR for incident CHD and stroke was elevated in the highest versus the lowest quartile of adherence (CHD: HR, 2.00; 95% CI, 1.53–2.61⁸⁸; stroke risk: HR, 1.39; 95% CI, 1.05–1.84).⁸⁷ Adjustment for this diet score was associated with a 63% mediation of the magnitude of estimated increased risk of stroke in African Americans <65 years of age.^{87,88}

Physical Inactivity

Current physical activity recommendations state that adults should engage in 150 min/wk of moderate-intensity aerobic physical activity or 75 minutes of vig-

orous-intensity aerobic physical activity and ≥ 2 d/wk of muscle-strengthening activities.⁸⁹ Data suggest that adherence to the recommended activity levels is particularly low among African Americans.⁹⁰ Fewer than 5% of all adults engaged in 30 minutes of moderate-intensity physical activity on most days of the week. Data from the CARDIA cohort show that more than one third of African American men and women report watching ≥ 4 hours of television per week, and this behavior is inversely associated with physical activity.⁹¹

When accelerometry is used to assess physical activity levels and sedentary behavior, patterns by race/ethnicity are less clear. In the NHANES study, there were no differences in minutes per day of moderate to vigorous activity between African Americans and whites.⁹⁰ Similarly, there was no difference in objectively determined sedentary behavior between African American and white men and women.⁹² In the REGARDS study, only 20% of African American men and 12% of women achieved >150 min/wk of accelerometer-determined moderate to vigorous physical activity per week compared with 30% and 20% of their white counterparts ($P<0.05$).⁹³ African Americans also spent statistically significantly more time engaged in sedentary behaviors as assessed by accelerometry in REGARDS (735 \pm 3 minutes for African American men versus 719 \pm 2 minutes for white men; 741 \pm 2 minutes for African American women versus 730 \pm 2 minutes for white women).⁹³

Cultural norms that influence behaviors, beliefs, and attitudes about physical activity are notable barriers to the adherence to physical activity recommendations. Some of these barriers include the perception that physical activity is “work” and not desired given the manual nature of daily jobs,⁹⁴ the lack of consistency between certain types of physical activity and African American self-identity (eg, double-dutch jump rope versus skiing),⁹⁵ and a low desirability for certain activities (eg, water sports) because of the physical nature,⁹⁶ the level of exertion or preparation required,^{97,98} and concerns about hairstyles, particularly among women.⁹⁹ Income is another proposed barrier to meeting physical activity guidelines,¹⁰⁰ but the role of income varies by geography in that it is a greater concern among rural adults¹⁰⁰ and less so among urban African Americans.¹⁰¹ Interpersonal barriers (eg, childcare, other family care), concerns about neighborhood safety, lack of access to facilities, and weather are also notable.⁹⁹ A significant limitation of the research on barriers to activity is that the vast majority has been conducted in African American women, and it is not known whether the same concerns are a priority among African American men.

Cigarette Smoking

Cigarette smoking is a strong and consistent risk factor for all CVDs, with little evidence of a differential magnitude of effect in African Americans and whites.¹⁰² Since

the height of smoking behavior during the 1960s, rates have declined across all racial/ethnic groups to current levels of 25.4% in African Americans and 25.8% in whites.¹⁰³ However, there are important age-related differences in the prevalence of cigarette smoking by race. Although cigarette smoking is markedly higher in white than African American adolescents (18.6% versus 8.2%, respectively),¹⁰⁴ these differences shrink considerably among adults >25 years of age.

Two primary areas of smoking-related disparities are exposure to environmental tobacco smoke (ie, secondhand smoke) and lower quit rates among African Americans. Environmental tobacco smoke is an established cardiovascular risk factor.¹⁰⁵ African American members of a health practice plan reported more environmental tobacco smoke exposure than whites,¹⁰⁶ and African American respondents to the NHIS (National Health Interview Survey)¹⁰⁷ and NHANES III¹⁰⁸ reported more environmental tobacco exposure. One hypothesis for the lower quit rates among African Americans than whites is that they are more likely to use menthol smoking products, which enhance the addictive potential of nicotine.¹⁰⁹ Tobacco companies target the marketing of mentholated products to African Americans (and youth),¹¹⁰ and among those smokers switching from mentholated to nonmentholated products, African Americans are more likely to revert back to the use of mentholated products.¹¹¹ Data from the Tobacco Use Supplement to the Current Population Survey report greater use of mentholated products among African American smokers (71%; 95% CI, 70.4–73.2) than white (21.0%; 95% CI, 20.5–21.4) or Hispanic (28.1%; 95% CI, 26.6–29.7) smokers. Smokers of mentholated products (of all races/ethnicities) were less likely to have quit.¹¹²

Clinical Sleep Disorders, Insufficient Sleep, and Poor-Quality Sleep

The relevance of sleep quality and duration on cardiovascular health was summarized in a 2016 AHA scientific statement.¹¹³ A growing body of research describes racial/ethnic disparities in sleep-disordered breathing, sleep duration, and sleep quality. Sleep has been hypothesized as a contributing factor to disparities in CVDs.¹¹⁴ African Americans are more likely to have obstructive sleep apnea (secondary to obesity),¹¹⁵ and untreated sleep apnea is associated with higher rates of cardiovascular mortality from CHD or stroke directly (eg, oxygen deprivation, sympathetic overreactivity, or inflammation¹¹⁶) or through the onset of other CVD risk factors such as hypertension or diabetes mellitus.¹¹⁷

Sleep duration is associated with cardiovascular risk factors and all-cause mortality; both short and long sleepers experience higher rates of events than those who sleep on average for 7 to 9 hours per night.^{118,119} African American respondents to the NHIS were 41%

more likely to self-report being short sleepers and 62% more likely to report being long sleepers than white participants.¹²⁰ African American participants reported longer sleep latency (time to fall asleep) and other adverse sleep symptoms than white and Hispanic participants.¹²¹ In the Chicago Area Sleep Study, a population-based epidemiological study of adults who underwent objective assessment of sleep via wrist actigraphy, the average sleep duration among African Americans was statistically significantly shorter per night (by ≈48 minutes) than among whites.¹²² A later report in the same cohort estimated that 11% of the disparity in hypertension prevalence between African Americans and whites was attributable to poor sleep quality (as determined by the percent of time during the sleep interval spent sleeping).¹²³

Multiple factors have been studied in relation to both short and long sleep duration in the all-African American JHS. Lower levels of education are associated with a greater likelihood of long sleep duration (OR, 2.19; 95% CI, 1.42–3.38). Similar findings were reported for the association of income with long sleep duration. Notably, individuals living in neighborhoods where they reported more neighborhood violence had shorter sleep duration (–9.82 minutes; 95% CI, –16.98 to –2.66) and poorer sleep quality.¹²⁴ In a later report, African American participants who reported higher levels of long-term stress were more likely to have short sleep duration (OR, 2.87; 95% CI, 2.02–4.08 versus the lower 3 quartiles of stress).¹²⁵

Less is known about racial differences in the impact of sleep duration and cardiovascular risk. Although long sleep duration appears to be a consistent risk factor for stroke, early indications suggest that among diabetic adults, short sleep duration (≤6 hours) is associated with increased stroke risk in whites (OR, 1.38; 95% CI, 1.06–1.80) but not in African Americans (OR, 0.86; 95% CI, 0.58–1.26).¹²⁶ Additional research is needed to test the association of objectively determined sleep duration with incident CHD risk in African Americans compared with whites.

Summary

There are significant disparities in the age of onset and prevalence of established CVD risk factors in African Americans. An earlier age of onset of obesity, hypertension, and diabetes mellitus is likely to contribute to the higher prevalence of these conditions and of CVD morbidity and mortality and to the lower life expectancy for African Americans versus whites. Adverse health behaviors in the African American population identified in the literature may explain in part the higher burden of CVD risk factors. Effective implementation of evidence-based guidelines could improve cardiovascular health and lower vascular risk in African Ameri-

cans. Ingrained cultural preferences and attitudes, as well as the social and physical context surrounding African Americans, influence the maintenance of behavior changes. Furthermore, experience shows that interventions targeting individuals are modestly successful during the period of intervention, but changes are not sustained beyond the intervention period. Consequently, population-wide strategies to influence health behavior change, as described in the AHA Community Guide for Prevention,¹²⁷ may have greater potential to reach both African American women and men and to shift the health behaviors and consequent cardiovascular risk of the entire population.

COMORBIDITIES

Certain health conditions that predispose to CVDs are more common among African Americans than whites. Below, we highlight chronic kidney disease, sickle cell disease/sickle cell trait, and HIV, given their relatively higher prevalence among African Americans.

Renal Disease (Chronic Kidney Disease and End-Stage Renal Disease)

The prevalence of chronic kidney disease, determined as estimated glomerular filtration rate $<60 \text{ mL}\cdot\text{min}^{-1}\cdot 1.73 \text{ m}^{-2}$ or albuminuria,¹²⁸ has increased in the United States according to NHANES data.^{129,130} African Americans have an excess burden of chronic kidney disease, resulting in part from the high prevalence of hypertension and diabetes mellitus,^{131,132} but that may also be the result of the percent of African admixture and other genetic factors.^{131,133,134} One such factor that has a higher prevalence in African Americans is sickle cell trait (discussed below), which has been associated with a higher rate of albuminuria and chronic kidney disease. Sickle cell trait carrier status was associated with 1.79 times higher (95% CI, 1.45–2.20) likelihood of developing chronic kidney disease over follow-up.¹³⁵

In a pooled analysis of the ARIC and CHS studies, the risk factor–adjusted HR for all-cause mortality in African Americans with versus without chronic kidney disease was 1.76 (95% CI, 1.35–2.31), whereas the HR for the same comparison among whites was significantly smaller (HR, 1.13; 95% CI, 1.02–1.26).¹³⁶ In the REGARDS study, African Americans with mild chronic kidney disease ($10 \leq \text{albumin/creatinine ratio} < 30$) versus no chronic kidney disease (albumin/creatinine ratio < 10) were more likely to experience CHD mortality (HR, 1.84; 95% CI, 1.34–2.53) than whites (HR, 1.23; 95% CI, 0.96–1.59). The disparities grew with severity of chronic kidney disease, and among participants with severe chronic kidney disease (albumin/creatinine ratio ≥ 300), the HRs were 3.21 (95% CI, 2.02–5.09) in African Americans and 1.49 (95% CI, 0.80–2.76) in

whites.¹³⁷ There was also evidence of effect modification in the relationship of albumin/creatinine ratio with stroke risk by race in REGARDS. The albumin/creatinine ratio was not associated with stroke risk among whites ($P > 0.05$), but among African Americans, the HR for a stroke event was 1.41 (95% CI, 1.01–1.96) for those with mild chronic kidney disease, 2.10 (95% CI, 1.48–2.99) for those with moderate chronic kidney disease, and 2.70 (95% CI, 1.58–4.61) for those with severe chronic kidney disease.¹³⁸

There are a number of paradoxical associations related to chronic kidney disease and end-stage renal disease among African Americans. The likelihood of progressing from chronic kidney disease to end-stage renal disease is greater in African Americans than whites.¹³⁹ Almost one third (32%) of the patients with end-stage renal disease are African American.¹⁴⁰ However, once on dialysis, survival is better among African Americans compared with whites. In 2008, the mortality rate for patients on dialysis was 16% in African Americans compared with 24% in whites.¹⁴⁰ Better survival on dialysis stands in contrast to the shorter life span for predialysis African Americans.¹⁴¹ A recent study also explored inflammation as a possible explanation for this observed paradox and found that this racial difference in end-stage renal disease survival did not exist at low levels of inflammation but is present at higher levels of inflammation as reflected by the upper tertile of C-reactive protein (CRP; $> 9.6 \text{ mg/L}$).¹⁴²

Sickle Cell Disease and Sickle Cell Trait

Sickle cell disease is a recessively inherited genetic condition caused by a glutamic acid to valine mutation in position 6 of the β -globin chain resulting in hemoglobin S formation. Subsequent sickling of red blood cells results from hemoglobin S–induced β -chain polymerization of hemoglobin tetramers.¹⁴³ Approximately 100 000 Americans have sickle cell disease, accounting for 1 in every 365 black births. However, 8% to 12% of African Americans carry a single mutation and have sickle cell trait. Although pulmonary complications and associated acute chest syndromes are the primary cardiopulmonary manifestations of sickle cell disease, there are vascular, thrombotic, metabolic, and myocardial complications of sickle cell disease and sickle cell trait.¹⁴⁴

Sudden death among patients with sickle cell disease is relatively high, with an incidence as high as 41% in some studies.¹⁴⁵ Patients with sickle cell disease have a variety of electrocardiographic abnormalities associated with malignant electric problems, including prolonged QTc, which may predispose to arrhythmias.¹⁴⁶ However, published reports of continuous electrocardiographic recordings in patients with sickle cell disease remain rare, and as a result, the effect of cardiac rhythm on

mortality in these patients is unknown. Emerging research also suggests that adults with sickle cell disease may have an imbalance of autonomic function, with a relative excess of sympathetic tone that might affect the initiation and progression of vaso-occlusive crisis, reflected by a propensity for arrhythmias, impaired cardiac perfusion, and an adverse hemodynamic profile.¹⁴⁷

Less is known about the cardiovascular risks for African Americans with sickle cell trait. There were no differences in the onset of CVD risk factors in a comparison of African American participants with and without sickle cell trait in the CARDIA study.¹⁴⁸ However, sickle cell trait was associated with an increased incidence of ischemic stroke in the ARIC study (HR, 1.4; 95% CI, 1.0–2.0).¹⁴⁹ In addition, emerging data suggest that African American athletes with sickle cell trait are at high risk for sudden death.¹⁵⁰ In a meta-analysis of the REGARDS, ARIC, MESA, JHS, and WHI (Women's Health Initiative) studies, sickle cell trait status was not significantly associated with incident myocardial infarction (HR, 1.10; 95% CI, 0.73–1.64) but was associated with CHD (HR, 1.42; 95% CI, 1.02–1.98) in a comparison of those with and without sickle cell trait.¹⁵¹ Further research on mechanisms and pathways is needed to explain the discrepancy between the 2 related outcomes.

HIV/AIDS

One of the emerging triumphs of 20th century medicine is the transformation of HIV/AIDS infection from a fatal condition to a chronic illness largely attributable to the introduction of highly active antiretroviral therapy. As a result, an increasing number of individuals with HIV infection are living longer and developing various cardiovascular conditions attributable to medications and the underlying inflammatory pathophysiology of HIV. A 2008 scientific statement¹⁵² presented evidence for the association of HIV with CVDs. Unfortunately, African Americans remain disproportionately affected by HIV infection, accounting for ~40% of the 1.2 million individuals living with HIV in the United States, and have the highest prevalence compared with other racial/ethnic groups.¹⁵³

A few studies include enough African American and white patients with HIV to evaluate the possibility of a differential effect of HIV on cardiovascular outcomes. In a systemic review of African Americans with HIV/AIDS in the United States, 2 of 5 studies indicated that African Americans with HIV were at increased CVD risk.¹⁵⁴ One of the few large assessments of CVD in African Americans with HIV is derived from a retrospective analysis of US National Hospital Discharge Surveys (1996–2008) showing that of 1.5 million discharges, the likelihood of hospitalization for CVD conditions was almost 50% higher (OR, 1.45; 95% CI, 1.39–1.51) in African Americans than in whites.¹⁵⁵ However, an unexpected

observation that warrants further investigation is that, when the subclinical coronary artery disease burden was compared between African American men with and without HIV in MACS (Multicenter AIDS Cohort Study), HIV positivity was associated with lower volume of total plaque.¹⁵⁶

Pharmacotherapy to manage HIV may increase the risks of CVDs. Although race-specific data were not presented, HIV-positive adults tend to have more dyslipidemia, characterized as high triglycerides and low HDL cholesterol (HDL-C) levels, likely as a result of decreased cholesterol ester transfer protein activity¹⁵⁷ compared with HIV-negative individuals. In the PURE study (Prospective Urban and Rural Epidemiology), patients with CD4 counts ≤ 200 cells/mm³ who were treated with nucleoside reverse transcriptase inhibitors (stavudine and lamivudine) and a nonnucleoside reverse transcriptase inhibitor (efavirenz or nevirapine) had significantly higher SBP, pulse pressure, and hemoglobin A_{1c} and more dyslipidemia over 5 years than untreated patients.¹⁵⁸

Summary

Although the excess burden of chronic kidney disease and end-stage renal disease in African Americans has origins in the disparate burden of cardiovascular risk factors, some unexpected observations are presented as related to faster rates of progression to end-stage renal disease but better survival on dialysis. It is possible that exploration of the healthcare experience of African Americans treated with dialysis could yield insights to provide better management for whites. Investigation of the health implications of sickle cell trait is ongoing, and with the availability of inexpensive genotyping for clinical and research purposes, we can investigate patterns in longitudinal studies in the population. As more research is carried out in diverse population cohorts of adults with HIV, we can gain additional insights into any disparities in the relationship of HIV status with CVDs in African Americans compared with other racial/ethnic groups.

CONTRIBUTION OF GENETICS TO DISPARITIES

As a multifactorial disease, CHD has both environmental and genetic underpinnings, with almost 300 variables identified to interact in unpredictable ways.¹⁵⁹ Conceptually, embracing the idea that genes segregate to populations is the most accurate way to think about genetics in general with respect to multifactorial diseases, including CVD. In studies genotyping populations, correlations have been made between populations and cardiovascular biomarkers, including inflammation, thrombosis, hypertension, lipid profiles, arrhythmia, and cardiac phenotype/left ventricular mass.

Genetic Loci for Inflammation

CRP is a heritable biomarker of systemic inflammation and predictor of CVD at the population level. Recent studies have performed genome-wide association studies (GWASs) of African American populations for genetic relationships to elevated CRP.^{160,161} In a GWAS of 8280 African American and 3548 Hispanic American postmenopausal women from the NHLBI SHARe (Single Nucleotide Polymorphism Health Association Resource), a unique triggering receptor expressed on myeloid cells 2 variant was associated with CRP in US minority populations, but genomic loci previously associated with CRP through GWASs of European populations demonstrated consistent patterns of association with CRP in African American and Hispanic American women.¹⁶²

In additional studies from the CARE study (Candidate Gene Association Resource) and race-combined meta-analyses of 29939 individuals of European descent, 4 loci were identified, 3 of which had been reported previously in populations of European descent.¹⁶⁰ Among African Americans, the fourth locus was the CD36 functional variant rs3211938, which is an extremely rare variant found in those of European descent. These findings were replicated in an independent sample of 8041 African Americans from WHI. In the race-combined meta-analyses, 13 loci reached significance, including 10 previously associated with CRP and 1 previously nominally associated with CRP.¹⁶⁰

Other studies that have investigated genes associated with elevated CRP levels in diverse populations have found similar gene associations.¹⁶⁰ In an investigation of 3109 African American and 6050 European Americans from the NHLBI ESP (Exome Sequencing Project) and CHARGE (Cohorts for Heart and Aging Research in Genomic Epidemiology) consortia, single-variant tests across candidate loci found an association of *APOE* ε2 rs7214 and higher CRP levels in African Americans.¹⁶³ Exome-wide, associations of *HNF1A*, *CRP*, *IL6R*, and *TOMM40-APOE* were confirmed.¹⁶³ Although these studies elucidated primarily genes overlapping with other populations, they also elucidated a few novel genes, demonstrating both common and unique potential contributors to elevated CRP levels in African Americans.

Genetic Loci for Thrombosis

Fibrinogen is a major component in the formation of a thrombus, cleaved by thrombin to form fibrin, the most abundant protein present in a blood clot.¹⁶⁴ Higher plasma fibrinogen levels are established markers of coronary artery disease, stroke, peripheral vascular disease, and atrial fibrillation.^{165–167} The heritability of plasma fibrinogen concentration as a predictor of CVD has been estimated to be 34% to 50%, with genetic variants so far explaining a small part (<2%) of this variation.¹⁶⁸

Fibrinogen levels have been reported to differ among European Americans, African Americans, and Africans, with the increased levels observed in those of African ancestry.^{169–173}

Evidence that genetic variants in the fibrinogen gene itself are related to cardiovascular risk was first identified in a GWAS of 6 population-based studies.¹⁷⁴ Since that time, a meta-analysis of 28 GWASs (which included 8289 African American participants) evaluating clinical outcomes has been published.¹⁶⁸ Twenty-four genome-wide independent signals from 23 loci were significant ($P < 5 \times 10^{-8}$), which included 15 novel associations that accounted for 3.7% of the plasma fibrinogen variation seen.¹⁷⁴ Enrichment analysis of these novel associations identified roles in fibrinogen regulation for the 3 structural genes and pathways related to inflammation, adipocytokines, and thyrotropin-releasing hormone signaling.¹⁶⁸ Although single-nucleotide polymorphisms (SNPs) in a few loci were significantly associated with coronary artery disease, the combined effect of the 24 fibrinogen-associated lead SNPs was not significant for coronary artery disease, stroke, or venous thromboembolism.¹⁶⁸

The fibrinogen γ chain has 2 splice variants, γ A and γ' , resulting in a poly-A signal in intron 9.¹⁷⁵ Because 8% to 15% of total fibrinogen is made of γ' fibrinogen, the variability of total fibrinogen and fibrinogen γ' in Africans is only partly explained by known CVD risk factors, with CRP being a major contributor.^{173,176} Fibrinogen SNPs account for 1.4% to 3.8% of the variance in total fibrinogen in African Americans^{177,178} and 2% of the variance in total fibrinogen in non-Hispanic blacks.¹⁷⁹ Recent studies have investigated the effect of fibrinogen and factor XIII genes on total and γ' fibrinogen and clot properties in black Africans.¹⁸⁰ Associations among total fibrinogen γ' levels, rs1049636 (fibrinogen γ chain), and rs2070011 (fibrinogen γ A promoter region) were identified.¹⁸⁰ SNPs interacted with total and/or γ' fibrinogen levels and clot properties in opposite ways, indicating that functionality should be a consideration in determining the effects of SNPs in CVD mechanisms and considerations of risk.¹⁸⁰

Genetic Loci for Hypertension

The heritability of hypertension documented in adoption, twin, and family studies suggests that 15% to 35% of the correlation may be genetic^{181–185}; hypertension onset before the age of 55 occurs nearly 4 times more frequently in individuals with a family history.¹⁸⁶ Together, the heritability of hypertension and increased predominance in African American populations suggest that unique genetic associations may be present in the development of CVD. Although studies of BP associations with cardiovascular phenotypes in European descendants are extensive,^{187,188} relatively few studies

have attempted to replicate these observations in African Americans.

In the first GWAS among African Americans, 80 000 SNPs in a discovery sample of 1017 African Americans from the Washington, DC, area were studied.¹⁸⁹ Multiple SNPs in genes encoding a Na⁺/K⁺/Ca²⁺ exchanger and voltage-dependent calcium channel, respectively, are significant genome-wide for SBP. No gene variants reached significance for association with diastolic BP or with hypertension as a binary trait.¹⁸⁹

Subsequent genome-wide and candidate gene association studies with SBP and diastolic BP (DBP) using the CARE consortium (consisting of 8591 African Americans) have identified additional genes related to DBP and SBP using 2 different genotyping platforms. None of these variants, however, were replicated in additional African American or European American cohorts, but 3 previously identified European American SNPs did replicate.¹⁹⁰ These findings support the notion that BP among African Americans has genetic influences on SBP and DBP at genetic loci found in European Americans, with potentially unique genes needing validation in other African American populations.¹⁹⁰

Admixture mapping is a method that can be used to detect disease variants with increased allele frequency differences in ancestral populations. Admixing mapping for SBP and DBP followed by trait marker associated in 6303 unrelated African American participants of the CARE consortium identified 5 significant genomic regions harboring genetic variants contributing to inter-individual BP variation.¹⁹¹ Overall, 3 loci were significantly associated with SBP and 1 with DBP and replicated in multiple large, independent studies of African Americans, including the WHI, Maywood, GENOA (Genetic Epidemiology Network of Arteriopathy), and HUFS (Howard University Family Study), and 1 native African sample (total replication size, 11 882).¹⁹¹ A novel variant on chromosome 5 (rs7726475) between the *SUB1* and *NPR3* genes was associated with SBP and DBP in the meta-analysis of the replication set.¹⁹¹ Meta-analyses of the CARE samples with the replication data identified a significant association of rs7726475 and DBP, highlighting the identification of genetic variants missed by GWASs.¹⁹¹

In an attempt to replicate the initial GWAS in African Americans described above, an independent sample of 2474 unrelated African Americans in the Milwaukee area (53% women, 47% men) was evaluated.¹⁹² When the top 16 associated SNPs plus the 8 SNPs associated with SBP and DBP in 2 genes (*STK-39* and *CDH-13*) found in European and Amish populations were investigated for their relationship with elevated BP in this African American cohort, no statistically significant differences were identified in African Americans, highlighting the importance of replication studies to validate the findings of GWASs.¹⁹²

Animal and human studies have demonstrated that genetic variations exist, and recent investigations have identified that rare polymorphisms in PCSK9 (proprotein convertase subtilisin/kexin type 9) are associated with BP in African American populations at high risk for CVD.¹⁹³ In an analysis of genomic data from the HyperGEN (Hypertension Genetic Epidemiology Network), 2 GWAS SNPs were identified with DBP (rs12048828: $\beta=1.8$, $P=0.05$; rs9730100: $\beta=1.0$, $P=0.05$) but were not significant after correction for multiple testing.¹⁹³ Although the DBP did not replicate, an association with SBP ($P=0.04$) did replicate in REGARDS, suggesting that rare variants in PCSK9 may influence BP among African Americans,¹⁹³ laying the groundwork for further validation studies and potential therapeutic considerations given the US Food and Drug Administration progress in approving PCSK9 inhibitors. Finally, in a discovery and meta-analysis that included 21 503 African Americans from across 16 studies, exome-centric single-variant and gene-based tests identified 31 new loci and 3 new genes associated with BP. Notably, these loci are enriched for known variants for other cardiometabolic traits, including dyslipidemia, inflammation, and insulin resistance.¹⁹⁴

Genetic Loci for Lipid Disorders

The initial GWASs investigating associations of genetic loci with LDL-C, HDL-C, and triglycerides in European ancestry populations identified 19 loci. To expand these associations with circulating lipid levels and CVD, index SNPs were genotyped at 19 loci in NHANES III ($n=7159$).¹⁹⁵ Analysis of non-Hispanic African Americans, Mexican Americans, and non-Hispanic whites identified the index SNP at 5 loci associated with LDL-C, HDL-C, or triglycerides in all 3 ethnic groups, which allowed the loci to be more finely mapped.¹⁹⁵

These subsequent studies determined that 22 SNPs in 13 candidate genes were associated with HDL-C, LDL-C, total cholesterol, and triglycerides.¹⁹⁶ Variants in *APOE* (rs7412, rs429358), *PON1* (rs854560), *ITGB3* (rs5918), and *NOS3* (rs2070744) associated with ≥ 1 of these lipids levels in at least 1 racial/ethnic population were found.¹⁹⁶ Multivariate linear regression analysis of 57 GWAS-identified or well-established lipid-related genetic loci with plasma concentrations of HDL-C, LDL-C, total cholesterol, triglycerides, total cholesterol/HDL-C ratio, and non-HDL-C was performed. With 1 exception (rs3764261 in *CEPT*), single SNP associations and the cumulative effect of multiple SNPs on blood lipid levels varied significantly by race/ethnicity. The findings were consistent for allele frequencies for all of the 57 GWAS-identified or lipid-related genetic loci.¹⁹⁷

As part of the NHLBI ESP, which included 1652 African Americans from CARDIA, CHS, ARIC, MESA, and WHI, participants who were heterozygous for any of

the 4 mutations in the *APOC3* gene had plasma triglyceride levels that were 39% lower than those who did not. Those same mutations were associated with a lower likelihood of CHD in whites, but the analysis has not yet included African Americans.¹⁹⁸ Another risk allele associated with CVD incidence is *APOL1*. Participants from the JHS who had 2 *APOL1* risk alleles had a doubled (OR, 2.17; $P=9.3\times 10^{-4}$) risk for incident CVD compared with those without a risk allele. These findings were replicated in the WHI for a combined OR of 2.12.¹⁹⁹

Expanding on these initial studies, the Population Architecture Using Genomics and Epidemiology Study was established to determine GWAS-identified variants in diverse population studies. Across racial/ethnic groups, a majority of the 55 of 60 replicated genotype-phenotype associations for HDL-C, LDL-C, and triglycerides in European Americans generalized to African American (48%, 61%, and 57%).²⁰⁰ For associations that did not generalize, differences in allele frequencies, linkage disequilibrium, and differences in effect sizes may contribute to the differences observed and offer insight into how next associations studies are designed in the future.²⁰⁰

To determine the validity of recent GWAS identification of loci/SNPs associated with plasma total cholesterol, LDL-C, HDL-C, and triglycerides, replication studies have been performed in 3 epidemiological samples comprising US non-Hispanic whites, US Hispanics, and African blacks.²⁰¹ In African blacks, 7 SNPs were significantly associated with at least 1 lipid trait, and 2 SNPs were associated with >1 lipid trait. These studies demonstrate the mixed results found with these loci with respect to various populations, each with its own allele frequency and contributions to disease.²⁰¹

PCSK9 (encoded by the *PCSK9* gene) is a regulator of LDL receptors, and therapies inhibiting this enzyme have shown promise as a novel, effective therapy for hyperlipidemia. However, in an exome array conducted to genotype >200 000 low-frequency sequences in 14 330 individuals of African ancestry, 4 low-frequency variants in the *PCSK9* gene were identified that had large effects on HDL-C or triglycerides, but none of these were associated with risk for CHD.²⁰²

Genetic Loci for Vascular Structure and Arrhythmia Risk

GWASs have investigated cardiac structure and systolic function in African Americans recently in the CARE study.²⁰³ Across the 9 cardiac phenotypes, 4 genetic loci reached significance for left ventricular mass, left ventricular internal diastolic diameter, interventricular septal wall thickness, and ejection fraction.²⁰³ None of these were identified in the European ancestry consortium, revealing unique African American variants enriched for

3 signaling pathways involving sonic hedgehog signaling, β -adrenergic signaling, and the oncostatin M signaling pathway.²⁰³ These 3 well-characterized pathways in cardiac remodeling suggest potential mechanisms underlying cardiac mechanisms associated with cardiac disease and potential targets for individualized therapies in individuals with these gene variants.²⁰³

The *SCN5A* gene, which plays a role in cardiac conduction and repolarization, was studied in relation to QT prolongation in the JHS. A common variant in individuals of African ancestry is the *SCN5A*-1103Y allele, which was found in 15.4% of JHS participants. In the 2% of participants with hypokalemia, there was a statistically significant ($P<0.005$) interaction whereby the allele was associated with prolongation of the QT interval by 15.6 milliseconds ($P=0.02$). In contrast, the association was modest for those without hypokalemia (4.1 milliseconds for each additional copy). The pattern of association between *SCN5A*-1103Y carrier status and hypokalemia was also observed for shorter QRS duration and longer QT, QTc, JT, and JTc intervals, and the findings were more pronounced among those with hypokalemia. The potential for diuretic-induced hypokalemia warrants consideration given the relatively high carrier rate of the *SCN5A*-1103Y allele and its attendant risks for sudden cardiac death.²⁰⁴

Summary

Concerns about the potential for the scientific community to misuse genetic information has been a barrier for many African Americans to participate in genetic research.^{205,206} However, education efforts that emphasize the value of such information for improving the health of African Americans could overcome many of these barriers. To date, genetic consortia have been the primary source of information on the contribution of genetics to CVD risk. As more cohorts that include racial/ethnic minorities join these collaborative efforts, the prevalence of risk alleles in minority cohorts can be determined, as well as their relationship with incident CVD risks. Finally, if the goals of personalized medicine are realized, these genomic findings may be combined with phenotypic information to provide precise characterizations of individual risk for CVDs.

DISEASE MANAGEMENT AND PREVENTION

Declines in CVD mortality are estimated to be attributable to the combination of the prevention of cardiovascular risk factors and the application of evidence-based therapies.²⁰⁷ Below, we discuss the challenge of screening for disease in African Americans with subclinical disease imaging, the role of risk calculators for risk strati-

fication, and the evidence for tailored pharmacological management of disease.

Subclinical CVD

Despite a higher burden of traditional risk factors and adverse health behaviors among African Americans, the prevalence of coronary artery calcium (CAC) is typically lower than among whites. Consequently, the inclusion of CAC in risk prediction equations could yield underestimates in African Americans.²⁰⁸ In the CARDIA study, the prevalence of CAC was 5% in African American women and 11% in African American men compared with 5% in white women and 18% in white men in adults 33 to 45 years old.²⁰⁹ In MESA participants (mean age, 62 years at baseline), CAC prevalence was highest among white men (70%), followed by African American men (52%), white women (45%), and African American women (37%).²¹⁰

In contrast, African Americans have been reported to have higher common carotid intima-media thickness but comparable internal carotid intima-media thickness compared with whites. According to MESA, the mean common carotid intima-media thickness was 0.91 mm in African Americans compared with 0.87 mm in whites, whereas the mean internal carotid intima-media thickness was similar in African Americans (1.11 mm) and whites (1.13 mm).²¹¹ The strength of association between carotid intima-media thickness and CAC also varies by racial/ethnic group, with the weakest association in African Americans.²¹¹ Genetic predisposition is a possible explanation for weaker associations among African Americans. In MESA, there was a positive association of European ancestry with CAC and common carotid intima-media thickness in African Americans.²¹² However, these associations were not replicated among African Americans in the CARDIA and CHS studies.^{213,214}

Differences in endothelial dysfunction between African Americans and whites have been observed in some settings but not others. In older adult women, brachial artery flow-mediated dilation was lower in African American compared with white women.²¹⁵ Among young men, microvascular function (peak and baseline forearm blood flow) is lower in African American men compared with white women.²¹⁶ In addition, nitric oxide bioavailability was lower in African American compared with white patients who were free from CVD risk factors, indicating poorer endothelial function.²¹⁷

Risk Calculators and Stratification

Until the turn of the century, there were no risk prediction tools based on data drawn on or validated in samples containing large numbers of African Americans.¹² A working group convened by the NHLBI on CHD risk prediction³ demonstrated that the FHS (Framingham

Heart Study) risk equation performed reasonably well in predicting CHD outcomes among African American participants in the ARIC study using data from the baseline examination in the 1980s.²¹⁸ In response to concerns about the validity of risk equations developed in an era when the CVD risk burden was markedly different, FHS investigators published an updated global risk equation in 2008 that broadened the outcomes beyond CHD.⁴ However, the transportability of the equation from the nearly all-white, New England-based cohort to the African American at-risk population remained uncertain.²¹⁸ The Reynolds Score improved prediction of ASCVD events by including family history and high-sensitivity CRP measurements in the prediction model. However, similar concerns arose about the generalizability of the score beyond the predominately white and upper-socioeconomic-status populations from which they were developed.²¹⁹

Risk functions for stroke were developed in the 1990s from the FHS⁷⁶ and the largely white CHS study.⁷⁷ Common predictors of stroke risk across cohorts were BP levels and treatment, prevalent diabetes mellitus, current smoking, atrial fibrillation, left ventricular hypertrophy, and heart diseases. Notably, dyslipidemia is not included in either equation, and there is a substantially heavier weight placed on hypertension as a risk factor. The CHS risk function additionally included measures of physical function and frailty (eg, timed walk). The Framingham Stroke Risk Function demonstrated good discrimination of stroke risk for whites and African Americans in the REGARDS study. However, the risk function likely overestimates stroke risk in the white population because of temporal declines in stroke risk, moving whites out of calibration and moving the higher-risk African Americans into calibration over time.²²⁰ That the risk factors for coronary disease and stroke differ underlies the somewhat modest correlation between the Framingham stroke and Framingham coronary risk functions (Spearman $\rho=0.68$).²²¹ These differences may explain the absence of an association between age-adjusted stroke and CHD mortality rates at the state level (Spearman $\rho=0.04$).²²¹

The Pooled Equations published recently by the AHA/American College of Cardiology Working Group directly address several of these possible shortcomings. They are a central pillar for the 4 guidelines simultaneously published to address modern assessment and management of CVD risk²²² and represent the first update on these topics since the publication of the Institute of Medicine's landmark "Guidelines We Can Trust."^{222a} The guideline for the assessment of cardiovascular risk features a new risk algorithm developed with data pooled from cohorts that included unprecedentedly large numbers of African Americans. One primary enhancement is that the risk model broadened the set of outcomes of concern beyond CHD to include stroke, a significant

source of disparity in CVD between African Americans and whites.^{223–225} The risk algorithm developed from the new guideline data outperformed other risk scores for initial ASCVD events among African Americans (and others) and has gained wide acceptance as a useful, broadly generalizable tool.²²⁴ Its easy accessibility online and its inclusion as a decision-support device through electronic medical record systems enhance its clinical utility. A limitation is that heart failure, another significant source of disparity, is not included. Finally, no risk prediction models are useful for the prediction of silent myocardial infarction, which may occur substantially more frequently among African Americans.

Hence, despite possible shortcomings, the Pooled Equations represent the best effort to date to produce an ASCVD event prediction tool for African American and white adults and has been validated in independent populations such as REGARDS.²²⁶ When this tool is applied to the NHANES data from 2007 to 2010, impressive disparities emerge: In the low-risk category, only 1.4% of African American men have a 10-year CVD risk of <2.5% compared with 18% of white men, 36.5% of African American women, and 47.1% of white women.²²⁶ Frequencies in other risk categories are less disparate; however, more than half (≈54%) of African American men and more than a third (≈34%) of African American women who are without known disease have a 10-year risk of >7.5% for an initial ASCVD event (compared with 44% for white men and 22% for white women), a key cut point for determining the intensity of therapy and the consideration of pharmacological management of CVD risk factors (eg, the use of statins for dyslipidemias).²²⁶

Recently, a race-specific tool has been developed by JHS investigators²²⁷ using data from the JHS.²²⁸ The investigators developed a tiered approach to select an algorithm that combined optimal prediction performance with ease of application in the primary care setting. They also expanded the relevance of the model to African American populations by including heart failure and stroke among the outcomes. Marginal improvements over the calibrated FHS and the new Pooled Equations were achieved by combining classic risk factors, brain natriuretic peptide, and ankle-brachial index measures. Clinical judgement is most often tested in cases in which the 10-year risk as determined by modern risk equations is neither high (currently meaning >7.5%) nor low (eg, a young person with substantial lifetime risk but with near-term low risk using the Pooled Equations). In such cases, the JHS findings suggest that imaging (echocardiography for left ventricular systolic performance), brain natriuretic peptide, and ankle-brachial index may offer additional useful data.

The authors concluded that, although the JHS-derived equations performed well in predicting disease and marginally improving risk classification over pre-

existing equations, they did not offer a substantial improvement in risk prediction among African Americans. This finding underscores (1) the enduring predictive power of the classic risk factors; (2) the fact that we can have reasonable confidence in the Pooled Equations in current wide use for African American and white American populations; (3) that any risk prediction tool is not intended as a substitute for a careful clinical assessment and may not be applicable for every patient; and (4) that a quantum leap forward in CVD prediction for our diverse population awaits further advances in the tools of precision medicine to assess environmental, genetic, and other determinants of risk.

Risk Stratification After ASCVD Events

Few algorithms addressing secondary prevention have come into widespread use because of confounding from treatment. In the REGARDS study, most of the traditional risk factors (eg, hypertension, diabetes mellitus, smoking, atrial fibrillation, left ventricular hypertrophy) play a similar role in predicting incident and recurrent stroke; however, a history of heart disease was associated with an RR of 1.42 (95% CI, 1.20–1.67) for incident stroke but only a nonsignificant (RR, 1.09; 95% CI, 0.85–1.40) risk for recurrent stroke.²²⁹ The substantial ($P=0.0002$) age-by-race interaction for incident stroke (with HRs of ≈3.0 at age 45 but 1.0 for age 85) was absent ($P=0.99$) for recurrent stroke, and no difference could be detected between the risk of recurrent strokes in African Americans and whites.²²⁹

Tailored Pharmacological Therapy in African Americans

Guidelines for pharmacological management of CVD in African Americans do not differ from management in other racial/ethnic groups. However, there are 2 notable exceptions, heart failure and hypertension, in which African American patients may benefit from tailored treatment approaches.

Heart Failure

The American College of Cardiology Foundation and AHA recommend angiotensin-converting enzyme (ACE) inhibitors, β -blockers, and aldosterone antagonists as the standard care in heart failure. Digoxin (Lanoxin) and diuretics are also recommended as adjuncts to control symptoms. Studies conducted in African Americans have documented that this population may have different responses to these medications compared with whites.²³⁰

ACE inhibitors are recommended for patients with New York Heart Association class I, II, III, or IV heart failure and for patients with left ventricular systolic dysfunction.²³¹ Despite robust evidence of benefit for ACE inhibitors for reducing mortality, the SOLVD (Studies of Left Ventricular Dysfunction) prevention and treat-

ment trial reported a smaller response to ACE inhibitor therapy (enalapril) in African American compared with white patients who had left ventricular dysfunction. Although SBP and DBP were lower among white patients treated with enalapril and the risk of hospitalization for heart failure was reduced by 44%, there were no significant reductions in any of these metrics among African American patients. Neither racial group received significant survival benefits from Enalapril treatment.²³²

β -blockers are recommended for patients with New York Heart Association class I, II, III, or IV heart failure and for patients with symptomatic and asymptomatic left ventricular systolic dysfunction.²³¹ Results from BEST (Beta-Blocker Evaluation of Survival Trial) reported a lack of significant survival benefit in African Americans with advanced heart failure when treated with bucindolol (HR, 1.17; 95% CI, 0.89–1.53), whereas there was an evident survival benefit in non-African American patients (HR, 0.82; 95% CI, 0.70–0.96).²³³ However, a recent meta-analysis using race-stratified data from COPENICUS (Carvedilol Prospective Randomized Cumulative Survival), MERIT-HF (Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure), and the US Carvedilol Heart Failure Study found potential survival benefit from bisoprolol, metoprolol, or carvedilol for African American patients with heart failure (RR, 0.67; 95% CI, 0.38–1.16), although this finding is not statistically significant.²³⁴ The absence of significant results may be attributable to the smaller sample size of African Americans in these studies.

Hydralazine plus isosorbide dinitrates is recommended to treat African American patients with left ventricular systolic dysfunction and advanced heart failure (New York Heart Association class III or IV), in addition to β -blockers and ACE inhibitors.^{230,231} Data from V-HeFT (Vasodilator-Heart Failure Trial) I showed that, compared with placebo, hydralazine plus isosorbide dinitrates significantly reduced mortality in African Americans, whereas such survival benefit was not observed among white patients. In addition, results from the V-HeFT II suggested that white patients received significantly more survival benefit from enalapril compared with hydralazine plus isosorbide dinitrates, whereas this treatment difference was not observed in African American patients. From these results, A-HeFT (African-American Heart Failure Trial) was designed for African American patients with New York Heart Association class III or IV heart failure only. This trial was terminated early because of a significant 43% improvement in survival in the hydralazine plus isosorbide dinitrates treatment group. In addition, the treatment group had a 33% reduction in the rate of first hospitalizations for heart failure and 52% improvement in quality of life.²³⁵ Another potential adjuvant therapy for heart failure in African American patients is supplemental aldosterone antagonist, particularly among those with left ventricu-

lar dysfunction. In the Genetic Risk of Heart Failure in African Americans trial, African Americans who had a common (62%) genetic polymorphism in the aldosterone synthase gene (TT genotype in *CYP11B2*) had higher levels of aldosterone and worse survival in heart failure.²³⁶

It is tempting to conclude that different heart failure therapies should be recommended to African American patients. However, most of the current studies on heart failure therapies included only a limited number of African American patients. Thus, the lack of significant findings is potentially attributable to small sample size. Although further study is needed, standard heart failure therapies should be used in African American patients with heart failure.

Hypertension

Disparities in hypertension control among African Americans are a primary source of disparities in CVDs. Reasons posited for the poorer control of elevated BP among African Americans usually focus on patient-related factors (eg, adherence issues, dietary indiscretion²³⁷), provider behavior (eg, inertia or poor regimen choices^{238,239}), or blunted efficacy of some widely used drug classes (eg, ACE inhibitors) in African Americans.²⁴⁰ However, the social and cultural environment is equally likely to influence the uptake and sustainability of preventive interventions. By carrying out research to elucidate the complexity of each of these dimensions of hypertension management (ie, social and cultural environment, behavior, pharmacology, pharmacogenetics), we can determine how the overall environmental/biopsychosocial milieu impedes or improves hypertension care and control in African Americans.²⁴¹ The ultimate goal is to generate greater precision in our care of African Americans with elevated BP.

There is evidence that race-specific treatment guidelines may be warranted. The British Hypertension Society suggests race-specific guideline recommendations to treat individuals of African descent with diuretics and calcium channel blockers and avoiding treatment with ACE inhibitors and β -blockers because of suppression of the renin-angiotensin system.²⁴² In the United States, the Eighth Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure provides specific recommendations for hypertension treatment in African Americans.²⁴³ One such recommendation is that a calcium channel blocker or thiazide-type diuretic be used as initial therapy for African American hypertensive patients.²⁴³ Another that has been met with some controversy is the recommendation that the SBP threshold for diagnosis and treatment should be raised from 140 to 150 mmHg in adults ≥ 60 years of age. The Association of Black Cardiologists and the Working Group on Women's Cardiovascular Health cite concerns about these

criteria given the large proportion of African Americans and women affected by hypertension and the risks of end-organ damage resulting from higher levels of BP.²⁴⁴

ALLHAT (Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial) provided the most evidence to support different responses to antihypertensive regimens in African American hypertensive patients.²⁴⁵ In the ALLHAT trial, patients on chlorthalidone achieved better BP control than patients on other therapies. Namely, SBP was 1 mm Hg lower than in the amlodipine group and 2 mm Hg lower than in the lisinopril group. The lowest risks for outcomes were also observed among those on chlorthalidone. Patients using ACE inhibitors (lisinopril) had a greater risk for stroke (RR, 1.40; 95% CI, 1.17–1.68), combined CVD (RR, 1.19; 95% CI, 1.09–1.30), and heart failure (RR, 1.30; 95% CI, 1.10–1.54) compared with those on chlorthalidone. These treatment differences were far more pronounced in African Americans compared with whites in the trial and make a strong argument in favor of diuretics as an initial drug of choice for treating hypertension in African Americans.²⁴⁵

The ALLHAT study also found a higher risk of stroke in African American hypertensive patients treated with ACE inhibitors (lisinopril) compared with a calcium channel blocker (amlodipine; RR, 1.51; 95% CI, 1.22–1.86); this association was not observed in non-African Americans (RR, 1.07; 95% CI, 0.89–1.28).²⁴⁵ These findings are consistent with results from several other studies. Using pooled results from 30 randomized controlled BP trials from 1968 to 2003, a meta-analysis study showed that there was no evident benefit from ACE inhibitors in achieving DBP goals for African American hypertensive patients. This study also found that African American hypertensive patients did not benefit from 1 particular β -blocker, atenolol, in reducing SBP.^{240,246} Similarly, several studies from the Veterans Affairs cooperative provide additional evidence that African American patients respond better to diuretics and calcium channel blockers.^{240,247–249}

Although African Americans are shown to be less responsive to ACE inhibitors, ACE inhibitors may offer benefits for African Americans with hypertensive renal disease. In AASK (African American Study of Kidney Disease and Hypertension), which is a comparison study of an ACE inhibitor (ramipril), a calcium channel blocker (amlodipine), and a β -blocker (metoprolol), ACE inhibition showed superior outcomes in relation to renal disease progression. Compared with calcium channel blockers and β -blockers, ACE inhibitors further slowed renal disease progression in African Americans with hypertensive renal disease and proteinuria. ACE inhibitors also offered clinical benefit in the combined end points of glomerular filtration rate events, end-stage renal disease, and death for African American hypertensive patients with renal disease with or without proteinuria.²⁵⁰

Adjustments to systems of care characterized by evidence-driven protocols embedded in a team-based, registry-documented, simplified, and tightly orchestrated approach have resulted in control rates of 87.1% in a population that 10 years earlier had a control rate of 43.6%.^{251,252} Until truly precise and personalized approaches are developed, a reliance on systems-level approaches may offer huge benefits. These approaches are being rapidly adopted by major caregiving organizations and have gained the endorsement of the AHA and the Centers for Disease Control and Prevention/Department of Health and Human Services–Sponsored Million Hearts Program.²⁵³

Summary

There is ample evidence that the traditional risk factors for CVD predict clinical outcomes equally well in African Americans and whites. More recent risk prediction equations that include stroke risk are particularly useful for African Americans. The utility of these equations for counseling about disease risks, and ultimately for prevention, can be enhanced when they are coupled with the selection of pharmacotherapies specifically recommended for disease prevention among African Americans.

SOCIAL AND CULTURAL ENVIRONMENT

The social and cultural environment in which African Americans live provides the context that influences the implementation of screening strategies, the application of risk prediction tools in clinical settings, and the uptake of evidence-based therapies.^{254–257} Existing research almost universally concludes that the combination of these factors complicates the prevention and management of CVD in African Americans. However, increased awareness and acknowledgement of these barriers have led to investment in strategies that work within the constraints of the environment to promote the cardiovascular health of African Americans.

In 2015, the AHA published a scientific statement addressing the social determinants of risk and outcomes of CVDs.²⁵⁸ The objective of that statement was to provide a comprehensive review of factors outside of biology that contribute to cardiovascular health and stand in the way of progress toward reaching the 2020 strategic Impact Goals.²⁵⁸ Unfortunately, African Americans face an overabundance of adverse social and environmental characteristics today in the United States.

African American race in the United States is closely correlated with socioeconomic class. Approximately 26% of African Americans are living in poverty compared with 13% of non-Hispanic whites and 15% of the overall population. The median family income is

\$43 151 in African American households compared with \$66 632 in the US population.²⁵⁹ On average, education levels are lower and health literacy is compromised. As a result, preventive health resources (healthy foods, safe spaces for physical activity, psychological stability resulting from occupational stability) are not as widely available to the majority of African Americans. Consequently, the challenge of improving health among African Americans requires a broader structural approach.

The AHA's community guide for improving health at the community level¹²⁷ discusses evidence-based strategies that can create environments that promote healthy behavior changes and support the health of community members. Given substantial evidence about the contribution of "built environment" factors such as limited access to supermarkets and healthier food choices,²⁶⁰ an overabundance of advertisements for high-calorie, low-nutrition foods and beverages,²⁶¹ and limited access to safe places for physical activity,²⁶² significant change may require investments in local community infrastructure, large-scale prevention programs, and social policies to support changes. These strategies include the use of policies to influence the built and social environments such as restricting the sale of nonnutritious foods in and around schools, labeling menus, providing incentives for food stores to build outlets in local food deserts, creating safe spaces for physical activity that are monitored to reduce the likelihood of crime, maintaining smoke-free restaurants and public spaces, and setting regulations about noise and air pollution from local businesses. These strategies may be particularly useful for closing the gaps in access and availability for African Americans and other minority communities who face more individual barriers to healthy behavior change and prevention.

There is considerable diversity within the African American community, with a large and growing middle- and upper-class community. However, despite higher education and more socioeconomic resources, health outcomes are still poorer in African Americans whose socioeconomic status is comparable to that of non-Hispanic whites.^{263,264} Historical factors, including housing laws and the migration patterns of racial/ethnic groups in the United States, inform the racial composition and structure of neighborhoods and communities. Research indicates that when African Americans live in predominately African American neighborhoods, health outcomes are worse.²⁶⁵ In the MESA study, the incidence of CVD went up 12% with every 1-SD increase in the degree of neighborhood segregation.²⁶⁶ Hypothesized reasons include variability in the availability of safe spaces for physical activity or access to healthy foods, which interfere with prevention. On a larger geographic scale, CVD event rates vary widely by geographic location.^{224,267} Nationally, 65% of Afri-

can Americans live in urban regions (counties classified by the National Center for Health Statistics as "large central metro" or "large fringe metro") compared with only 49% of non-Hispanic whites.²⁶⁸ Stroke mortality rates are 30% higher in rural regions of the country than in urban regions.²⁶⁹

Another explanation for persistent disparities across the socioeconomic range in African Americans is the multiple sources of stress and unique sources of stress faced by African Americans. Common sources of stress, job stress and strain, neighborhood-related safety concerns, socioeconomic concerns, and major life events combine with salient sources of stress for African Americans such as perceived discrimination.^{270–273} Perceived racial discrimination is known to be associated with all aspects of health, including hypertension,²⁷⁴ weight gain and obesity,²⁷⁵ persistent inflammation and other subclinical processes,^{276,277} and incident CVD events.²⁷⁸ The potential for interventions that promote positive psychological health (eg, mindfulness, resiliency) to reduce stress levels may be relevant in improving the health behaviors and ultimately health outcomes of African Americans.^{279,280}

The effectiveness of behavior change interventions in the African American community is compromised by a number of factors. As a result of ingrained cultural practices or fears surrounding the healthcare system because of historical abuses, African Americans may be less likely to follow physician recommendations for behavior change.²⁸¹ One of the most striking examples of the confluence of cultural practices, socioeconomic conditions, and attitudes among African Americans compromising the effectiveness of therapy is in management of the obesity epidemic. Although it is possible that lower levels of overall education or health literacy may contribute to the lack of understanding of the relationship among dietary intake, body weight, and chronic disease risk²⁸² or that food purchasing habits are influenced by socioeconomic circumstances,²⁵⁹ it is equally likely that traditional preferences may be more influential than socioeconomic status in predicting food purchasing behaviors.²⁸³ According to some reports, African American families and social networks do not promote positive lifestyle changes.^{213,284,285} The attitudes and behaviors that some African Americans have about weight may also influence the adoption of dietary guidelines.^{75,286–288}

As a result, although individual weight-loss interventions show variable effectiveness, all weight-loss interventions are systematically less effective in African Americans according to a meta-analysis.²⁸⁹ In a review of weight-loss interventions, studies that were successful included cultural adaptation and, even more important, constituent involvement; that is, adaptation informed by the experience of the target group.²⁹⁰ Conducting interventions in locations that are valued in the African

American community can enhance their uptake; 1 review reports that 70% of interventions conducted in African American faith-based organizations achieved success in weight reduction.²⁹¹

Summary

Although African Americans face a number of social and structural barriers to positive cardiovascular health, a number of strengths of the cultural environment can be leveraged to disseminate behavioral health interventions (eg, the central role of the church for reaching women and older adults). Some of these barriers can be addressed by targeting the macroenvironment via policy changes at the federal (eg, Affordable Care Act), state (eg, cigarette smoking bans), and local (eg, food availability in schools) levels. Doing so could create environments to support and sustain positive health behavior changes. However, finding strategies that reach younger African Americans and men with disease prevention messages is a challenge that must be met to change the trajectory of health in the African American community.

THE NEXT STEPS

To make progress toward our goal of promoting health equity and achieving the AHA 2020 Impact Goals, the significant burden of morbidity and mortality from CVD among African Americans must be reduced. Doing so will require collaborations across multiple disciplines, both within and outside of the traditional umbrella of healthcare providers given the complex historical, social, and economic reasons why African Americans experience poorer cardiovascular health. Efforts to carry out high-quality research studies should be supplemented by additional research on the dissemination and implementation of effective interventions to modify health behaviors and to mitigate CVD risks. New initiatives such as the precision medicine initiative (ie, All of Us) provide a unique opportunity to learn more about individual genotypes and how therapies can be tailored to be maximally effective for African Americans.

However, such research will be effective only if the diverse spectrum of the African American population is represented. Despite the wealth of information that we have about the CVD burden in African Americans from surveillance studies and longitudinal cohort studies, African American men and socioeconomically disenfranchised (including homeless) populations remain less likely to participate in research. Although these population subgroups are difficult to reach in any racial/ethnic group, historical abuses by the healthcare system (eg, Tuskegee Syphilis Study), which spurred a lasting culture of mistrust, may further alienate African Americans. At present, life expectancy among African American men is shorter than all for other racial/eth-

nic groups, and health outcomes among the socioeconomically disenfranchised in the United States mirror those of populations in the developing world. Identifying strategies to reach these groups with preventive and clinical care needs to be a high priority.

One strategy to begin to repair trust and to increase the engagement of all African Americans in the healthcare system is to diversify the workforce of healthcare professionals. In a recent report from an NHLBI Think Tank on Strategies to Promote Health Equity, a priority was placed on ensuring a diverse workforce of clinicians and researchers by creating unique transdisciplinary training programs.²⁷¹ From the research perspective, identifying and addressing the numerous barriers to cardiovascular health requires input from disciplines as broad as anthropology, public policy, and education, as well as traditional disciplines such as medicine, epidemiology, and psychology. On the clinical side, healthcare providers who demonstrate cultural competency in their interactions and a willingness to try to understand the perspectives of the patients they treat will yield higher-quality interactions. Although such skills are relevant for all healthcare providers, the infusion of the patient-facing medical fields with more African American clinicians (eg, physicians, physician assistants, and nursing professionals) may speed the uptake of these important principles. However, efforts to increase the number of clinical providers must begin with programming that builds the pipeline of African American students who are interested in selecting medicine as a career and who are prepared to succeed.

With the considerable amount of information that we have about the prevalence and sources of disparities in cardiovascular health between African Americans and other racial/ethnic groups, we stand poised to address and eliminate those disparities with contributions from professionals with expertise in basic science and pharmacology, clinical medicine, and public health. By successfully translating findings from across disciplines, scientists and practitioners from other disciplines can apply innovative strategies to improve the cardiovascular health of African Americans.

FOOTNOTES

The American Heart Association makes every effort to avoid any actual or potential conflicts of interest that may arise as a result of an outside relationship or a personal, professional, or business interest of a member of the writing panel. Specifically, all members of the writing group are required to complete and submit a disclosure questionnaire showing all such relationships that might be perceived as real or potential conflicts of interest.

This statement was approved by the American Heart Association Science Advisory and Coordinating Committee on June 23, 2017, and the American Heart Association Executive Committee on August 21, 2017. A copy of the document is available at <http://professional.heart.org/statements> by using

either "Search for Guidelines & Statements" or the "Browse By Topic" area. To purchase additional reprints, call 843-216-2533 or e-mail kelle.ramsay@wolterskluwer.com.

The American Heart Association requests that this document be cited as follows: Carnethon MR, Pu J, Howard G, Albert MA, Anderson CAM, Bertoni AG, Mujahid MS, Palaniappan L, Taylor HA Jr, Willis M, Yancy CW; on behalf of the American Heart Association Council on Epidemiology and Prevention; Council on Cardiovascular Disease in the Young; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Functional Genomics and Translational Biology; and Stroke Council. Cardiovascular health in African Americans: a scientific statement from the American Heart Association. *Circulation*. 2017;136:e393–e423. doi: 10.1161/CIR.0000000000000534.

Expert peer review of AHA Scientific Statements is conducted by the AHA Office of Science Operations. For more on AHA statements and guidelines development, visit <http://professional.heart.org/statements>. Select the "Guidelines & Statements" drop-down menu, then click "Publication Development."

Permissions: Multiple copies, modification, alteration, enhancement, and/or distribution of this document are not permitted without the express permission of the American Heart Association. Instructions for obtaining permission are located at http://www.heart.org/HEARTORG/General/Copyright-Permission-Guidelines_UCM_300404_Article.jsp. A link to the "Copyright Permissions Request Form" appears on the right side of the page.

Circulation is available at <http://circ.ahajournals.org>.

DISCLOSURES

Writing Group Disclosures

Writing Group Member	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory Board	Other
Mercedes R. Carnethon	Northwestern University Preventive Medicine	None	None	None	None	None	None	None
Michelle A. Albert	University of California, San Francisco	Kellogg Foundation (supports the study of minority populations, CVD, and adversity)*; NIH/NIA*	None	None	None	None	None	None
Cheryl A.M. Anderson	University of California at San Diego	None	None	None	None	None	None	None
Alain G. Bertoni	Wake Forest University School of Medicine, Epidemiology and Prevention Medical Center	NIH†	None	None	None	None	None	None
George Howard	University of Alabama at Birmingham School of Public Health	NIH†	None	None	None	None	Bayer Health Care*	None
Mahasin S. Mujahid	University of California, Berkeley	NIH/NHLBI K01HL115494†	None	None	None	None	Morehouse School of Medicine Cardiovascular Research Institute*	None
Latha Palaniappan	Stanford University	None	None	None	None	None	None	None
Jia Pu	Mathematica Policy Research Health Research	None	None	None	None	None	None	None
Herman A. Taylor Jr.	Morehouse School of Medicine	NIH†	None	None	None	None	None	None
Monte Willis	University of North Carolina Pathology and Laboratory Medicine, and McAllister Heart Institute	LeDucq Foundation†; NIH†	None	None	None	None	None	None
Clyde W. Yancy	Northwestern University Internal Medicine/ Cardiology	None	None	None	None	None	None	None

This table represents the relationships of writing group members that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all members of the writing group are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

†Significant.

Reviewer Disclosures

Reviewer	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory Board	Other
Adolfo Correa	University of Mississippi Medical Center	NHLBI (PI of the JHS) [†]	None	None	None	None	None	None
Gladys Velarde	University of Florida	Gilead Pharmaceutical*	None	None	None	None	None	None
Karol Watson	UCLA	None	None	None	None	None	None	None

This table represents the relationships of reviewers that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all reviewers are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

[†]Significant.

REFERENCES

- Rosamond WD, Chambless LE, Heiss G, Mosley TH, Coresh J, Whitsel E, Wagenknecht L, Ni H, Folsom AR. Twenty-two-year trends in incidence of myocardial infarction, coronary heart disease mortality, and case fatality in 4 US communities, 1987–2008. *Circulation*. 2012;125:1848–1857.
- Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, de Ferranti SD, Floyd J, Fornage M, Gillespie C, Isasi CR, Jimenez MC, Jordan LC, Judd SE, Lackland D, Lichtman JH, Lisabeth L, Liu S, Longenecker CT, Mackey RH, Matsushita K, Mozaffarian D, Mussolino ME, Nasir K, Neumar RW, Palaniappan L, Pandey DK, Thiagarajan RR, Reeves MJ, Ritchey M, Rodriguez CJ, Roth GA, Rosamond WD, Sasson C, Towfighi A, Tsao CW, Turner MB, Virani SS, Voeks JH, Willey JZ, Wilkins JT, Wu JHY, Alger HM, Wong SS, Muntner P; on behalf of the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2017 update: a report from the American Heart Association [published correction appears in *Circulation* 2017;136:e196]. *Circulation*. 2017;135:e146–e603. DOI: 10.1161/CIR.0000000000000485.
- Bonow RO, Grant AO, Jacobs AK. The cardiovascular state of the union: confronting healthcare disparities. *Circulation*. 2005;111:1205–1207. doi: 10.1161/01.CIR.0000160705.97642.92.
- Palaniappan LP, Araneta MR, Assimes TL, Barrett-Connor EL, Carnethon MR, Criqui MH, Fung GL, Narayan KM, Patel H, Taylor-Piliae RE, Wilson PW, Wong ND; on behalf of the American Heart Association Council on Epidemiology and Prevention; American Heart Association Council on Peripheral Vascular Disease; American Heart Association Council on Nutrition, Physical Activity, and Metabolism; American Heart Association Council on Clinical Cardiology; American Heart Association Council on Cardiovascular Nursing; Council on Cardiovascular Nursing. Call to action: cardiovascular disease in Asian Americans: a science advisory from the American Heart Association [published correction appears in *Circulation*. 2010;122:e516]. *Circulation*. 2010;122:1242–1252. doi: 10.1161/CIR.0b013e3181f22af4.
- Rodriguez CJ, Allison M, Daviglus ML, Isasi CR, Keller C, Leira EC, Palaniappan L, Piña IL, Ramirez SM, Rodriguez B, Sims M; on behalf of the American Heart Association Council on Epidemiology and Prevention; American Heart Association Council on Clinical Cardiology; American Heart Association Council on Cardiovascular and Stroke Nursing. Status of cardiovascular disease and stroke in Hispanics/Latinos in the United States: a science advisory from the American Heart Association. *Circulation*. 2014;130:593–625. doi: 10.1161/CIR.0000000000000071.
- US Census Bureau. National population projections: summary tables. 2012. <https://www.census.gov/data/tables/2012/demo/popproj/2012-summary-tables.html>. Accessed March 17, 2017.
- Arias E, Heron M, Xu JQ. United States life tables, 2012. *Natl Vital Stat Rep*. 2016;65:1–65.
- Gillespie CD, Wigington C, Hong Y; Centers for Disease Control and Prevention (CDC). Coronary heart disease and stroke deaths—United States, 2009. *MMWR Suppl*. 2013;62:157–160.
- Centers for Disease Control and Prevention (CDC). Prevalence of coronary heart disease—United States, 2006–2010. *MMWR Morb Mortal Wkly Rep*. 2011;60:1377–1381.
- Safford MM, Brown TM, Muntner PM, Durant RW, Glasser S, Halanynch JH, Shikany JM, Prineas RJ, Samdarshi T, Bittner VA, Lewis CE, Gamboa C, Cushman M, Howard V, Howard G; REGARDS Investigators. Association of race and sex with risk of incident acute coronary heart disease events. *JAMA*. 2012;308:1768–1774. doi: 10.1001/jama.2012.14306.
- Bibbins-Domingo K, Pletcher MJ, Lin F, Vittinghoff E, Gardin JM, Arynchyn A, Lewis CE, Williams OD, Hulley SB. Racial differences in incident heart failure among young adults. *N Engl J Med*. 2009;360:1179–1190. doi: 10.1056/NEJMoa0807265.
- Bahrami H, Kronmal R, Bluemke DA, Olson J, Shea S, Liu K, Burke GL, Lima JA. Differences in the incidence of congestive heart failure by ethnicity: the Multi-Ethnic Study of Atherosclerosis. *Arch Intern Med*. 2008;168:2138–2145. doi: 10.1001/archinte.168.19.2138.
- Loehr LR, Rosamond WD, Chang PP, Folsom AR, Chambless LE. Heart failure incidence and survival (from the Atherosclerosis Risk in Communities study). *Am J Cardiol*. 2008;101:1016–1022. doi: 10.1016/j.amjcard.2007.11.061.
- Zheng ZJ, Croft JB, Giles WH, Mensah GA. Sudden cardiac death in the United States, 1989 to 1998. *Circulation*. 2001;104:2158–2163.
- Chan PS, Nichol G, Krumholz HM, Spertus JA, Jones PG, Peterson ED, Rathore SS, Nallamothu BK, American Heart Association National Registry of Cardiopulmonary Resuscitation (NRCPR) Investigators. Racial differences in survival after in-hospital cardiac arrest. *JAMA*. 2009;302:1195–1201. doi: 10.1001/jama.2009.1340.
- Lackland DT, Rocella EJ, Deutsch AF, Fornage M, George MG, Howard G, Kissela BM, Kittner SJ, Lichtman JH, Lisabeth LD, Schwamm LH, Smith EE, Towfighi A; on behalf of the American Heart Association Stroke Council; Council on Cardiovascular and Stroke Nursing; Council on Quality of Care and Outcomes Research; Council on Functional Genomics and Translational Biology. Factors influencing the decline in stroke mortality: a statement from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:315–353. doi: 10.1161/01.str.0000437068.30550.cf.
- Rosamond WD, Folsom AR, Chambless LE, Wang CH, McGovern PG, Howard G, Copper LS, Shahar E. Stroke incidence and survival among middle-aged adults: 9-year follow-up of the Atherosclerosis Risk in Communities (ARIC) cohort. *Stroke*. 1999;30:736–743.
- Yang D, Howard G, Coffey CS, Roseman J. The confounding of race and geography: how much of the excess stroke mortality among African Americans is explained by geography? *Neuroepidemiology*. 2004;23:118–122. doi: 10.1159/000075954.
- Kleindorfer D, Broderick J, Khoury J, Flaherty M, Woo D, Alwell K, Moomaw CJ, Schneider A, Miller R, Shukla R, Kissela B. The unchanging incidence and case-fatality of stroke in the 1990s: a population-based study. *Stroke*. 2006;37:2473–2478. doi: 10.1161/01.STR.0000242766.65550.92.
- Broderick JP, Brott T, Tomsick T, Huster G, Miller R. The risk of subarachnoid and intracerebral hemorrhages in blacks as compared with whites. *N Engl J Med*. 1992;326:733–736. doi: 10.1056/NEJM199203123261103.
- Sturgeon JD, Folsom AR, Longstreth WT Jr, Shahar E, Rosamond WD, Cushman M. Risk factors for intracerebral hemorrhage in a pooled prospective study. *Stroke*. 2007;38:2718–2725. doi: 10.1161/STROKEAHA.107.487090.
- Howard G, Cushman M, Howard VJ, Kissela BM, Kleindorfer DO, Moy CS, Switzer J, Woo D. Risk factors for intracerebral hemorrhage: the REasons

- for Geographic And Racial Differences in Stroke (REGARDS) study [published correction appears in *Stroke*. 2013;44:e63]. *Stroke*. 2013;44:1282–1287. doi: 10.1161/STROKEAHA.111.000529.
23. Criqui MH, Aboyans V. Epidemiology of peripheral artery disease. *Circ Res*. 2015;116:1509–1526. doi: 10.1161/CIRCRESAHA.116.303849.
 24. Allison MA, Ho E, Denenberg JO, Langer RD, Newman AB, Fabsitz RR, Criqui MH. Ethnic-specific prevalence of peripheral arterial disease in the United States [published correction appears in *Circ Res*. 2015;117:e12]. *Am J Prev Med*. 2007;32:328–333. doi: 10.1016/j.amepre.2006.12.010.
 25. Newman AB, Siscovick DS, Manolio TA, Polak J, Fried LP, Borhani NO, Wolfson SK. Ankle-arm index as a marker of atherosclerosis in the Cardiovascular Health Study: Cardiovascular Heart Study (CHS) Collaborative Research Group. *Circulation*. 1993;88:837–845.
 26. Zheng ZJ, Sharrett AR, Chambless LE, Rosamond WD, Nieto FJ, Sheps DS, Dobs A, Evans GW, Heiss G. Associations of ankle-brachial index with clinical coronary heart disease, stroke and preclinical carotid and popliteal atherosclerosis: the Atherosclerosis Risk in Communities (ARIC) Study. *Atherosclerosis*. 1997;131:115–125.
 27. Allison MA, Criqui MH, McClelland RL, Scott JM, McDermott MM, Liu K, Folsom AR, Bertoni AG, Sharrett AR, Homma S, Kori S. The effect of novel cardiovascular risk factors on the ethnic-specific odds for peripheral arterial disease in the Multi-Ethnic Study of Atherosclerosis (MESA). *J Am Coll Cardiol*. 2006;48:1190–1197. doi: 10.1016/j.jacc.2006.05.049.
 28. Ix JH, Allison MA, Denenberg JO, Cushman M, Criqui MH. Novel cardiovascular risk factors do not completely explain the higher prevalence of peripheral arterial disease among African Americans: the San Diego Population Study. *J Am Coll Cardiol*. 2008;51:2347–2354. doi: 10.1016/j.jacc.2008.03.022.
 29. Deleted in proof.
 30. Ning H, Labarthe DR, Shay CM, Daniels SR, Hou L, Van Horn L, Lloyd-Jones DM. Status of cardiovascular health in US children up to 11 years of age: the National Health and Nutrition Examination Surveys 2003–2010. *Circ Cardiovasc Qual Outcomes*. 2015;8:164–171. doi: 10.1161/CIRCOUTCOMES.114.001274.
 31. Shay CM, Ning H, Daniels SR, Rooks CR, Gidding SS, Lloyd-Jones DM. Status of cardiovascular health in US adolescents: prevalence estimates from the National Health and Nutrition Examination Surveys (NHANES) 2005–2010. *Circulation*. 2013;127:1369–1376. doi: 10.1161/CIRCULATIONAHA.113.001559.
 32. National Center for Health Statistics. *Health, United States, 2015: With Special Feature on Racial and Ethnic Health Disparities*. Hyattsville, MD: National Center for Health Statistics; 2016. Report 2016-1232.
 33. Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, Chen J, He J. Global disparities of hypertension prevalence and control: a systematic analysis of population-based studies from 90 countries. *Circulation*. 2016;134:441–450. doi: 10.1161/CIRCULATIONAHA.115.018912.
 34. Marden JR, Walter S, Kaufman JS, Glymour MM. African ancestry, social factors, and hypertension among non-Hispanic blacks in the Health and Retirement Study. *Biodemography Soc Biol*. 2016;62:19–35. doi: 10.1080/19485565.2015.1108836.
 35. Non AL, Gravlee CC, Mulligan CJ. Education, genetic ancestry, and blood pressure in African Americans and whites. *Am J Public Health*. 2012;102:1559–1565. doi: 10.2105/AJPH.2011.300448.
 36. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics*. 2004;114(suppl 4th report):555–576.
 37. Bao W, Threefoot SA, Srinivasan SR, Berenson GS. Essential hypertension predicted by tracking of elevated blood pressure from childhood to adulthood: the Bogalusa Heart Study. *Am J Hypertens*. 1995;8:657–665. doi: 10.1016/0895-7061(95)00116-7.
 38. Carson AP, Howard G, Burke GL, Shea S, Levitan EB, Muntner P. Ethnic differences in hypertension incidence among middle-aged and older adults: the Multi-Ethnic Study of Atherosclerosis. *Hypertension*. 2011;57:1101–1107. doi: 10.1161/HYPERTENSIONAHA.110.168005.
 39. Howard G, Safford MM, Moy CS, Howard VJ, Kleindorfer DO, Unverzagt FW, Soliman EZ, Flaherty ML, McClure LA, Lackland DT, Wadley VG, Pulley LV, Cushman M. Racial differences in the incidence of cardiovascular risk factors in older black and white adults. *J Am Geriatr Soc*. 2017;65:83–90. doi: 10.1111/jgs.14472.
 40. Yoon SS, Burt V, Louis T, Carroll MD. Hypertension among adults in the United States, 2009–2010. *NCHS Data Brief*. 2012:1–8.
 41. Gu Q, Burt VL, Dillon CF, Yoon S. Trends in antihypertensive medication use and blood pressure control among United States adults with hypertension: the National Health and Nutrition Examination Survey, 2001 to 2010. *Circulation*. 2012;126:2105–2114. doi: 10.1161/CIRCULATIONAHA.112.096156.
 42. Howard G, Prineas R, Moy C, Cushman M, Kellum M, Temple E, Graham A, Howard V. Racial and geographic differences in awareness, treatment, and control of hypertension: the REasons for Geographic And Racial Differences in Stroke study. *Stroke*. 2006;37:1171–1178. doi: 10.1161/01.STR.0000217222.09978.ce.
 43. Howard G, Banach M, Cushman M, Goff DC, Howard VJ, Lackland DT, McVay J, Meschia JF, Muntner P, Oparil S, Rightmyer M, Taylor HA. Is blood pressure control for stroke prevention the correct goal? The lost opportunity of preventing hypertension. *Stroke*. 2015;46:1595–1600. doi: 10.1161/STROKEAHA.115.009128.
 44. Whelton PK, Einhorn PT, Muntner P, Appel LJ, Cushman WC, Diez Roux AV, Ferdinand KC, Rahman M, Taylor HA, Ard J, Arnett DK, Carter BL, Davis BR, Freedman BI, Cooper LA, Cooper R, Desvigne-Nickens P, Gavini N, Go AS, Hyman DJ, Kimmel PL, Margolis KL, Miller ER 3rd, Mills KT, Mensah GA, Navar AM, Ogedegbe G, Rakotz MK, Thomas G, Tobin JN, Wright JT, Yoon SS, Cutler JA; for the National Heart, Lung, and Blood Institute Working Group on Research Needs to Improve Hypertension Treatment and Control in African Americans. Research Needs to Improve Hypertension Treatment and Control in African Americans. *Hypertension*. 2016;68:1066–1072. doi: 10.1161/HYPERTENSIONAHA.116.07905.
 45. American Diabetes Association. 2. Classification and diagnosis of diabetes. *Diabetes Care*. 2016;39:S13–S22.
 46. Menke A, Casagrande S, Geiss L, Cowie CC. Prevalence of and trends in diabetes among adults in the United States, 1988–2012. *JAMA*. 2015;314:1021–1029. doi: 10.1001/jama.2015.10029.
 47. Selvin E, Parrinello CM, Sacks DB, Coresh J. Trends in prevalence and control of diabetes in the United States, 1988–1994 and 1999–2010. *Ann Intern Med*. 2014;160:517–525. doi: 10.7326/M13-2411.
 48. Writing Group for the SEARCH for Diabetes in Youth Study Group, Dabelea D, Bell RA, D'Agostino RB Jr, Imperatore G, Johansen JM, Linder B, Liu LL, Loots B, Marcovina S, Mayer-Davis EJ, Pettitt DJ, Waitzfelder B. Incidence of diabetes in youth in the United States [published correction appears in *JAMA*. 2007;298:627]. *JAMA*. 2007;297:2716–2724.
 49. Lacy ME, Wellenius GA, Sumner AE, Correa A, Carnethon MR, Liem RI, Wilson JG, Sacks DB, Jacobs DR Jr, Carson AP, Luo X, Gjelsvik A, Reiner AP, Naik RP, Liu S, Musani SK, Eaton CB, Wu WC. Association of sickle cell trait with hemoglobin A1c in African Americans. *JAMA*. 2017;317:507–515. doi: 10.1001/jama.2016.21035.
 50. Centers for Medicare & Medicaid Services. ACO shared savings program quality measures. 2017. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/sharedsavingsprogram/Downloads/MSSP-QM-Benchmarks-2016.pdf>. Accessed March 22, 2017.
 51. Golden SH, Brown A, Cauley JA, Chin MH, Gary-Webb TL, Kim C, Sosa JA, Sumner AE, Anton B. Health disparities in endocrine disorders: biological, clinical, and nonclinical factors: an Endocrine Society scientific statement. *J Clin Endocrinol Metab*. 2012;97:E1579–E1639. doi: 10.1210/jc.2012-2043.
 52. National Center for Health Statistics. Mortality data—vital statistics. NCHS' Multiple Cause of Death Data, 1959–2015. <http://www.nber.org/data/vital-statistics-mortality-data-multiple-cause-of-death.html>. Accessed March 17, 2017.
 53. Nsiah-Kumi P, Ortmeier SR, Brown AE. Disparities in diabetic retinopathy screening and disease for racial and ethnic minority populations: a literature review. *J Natl Med Assoc*. 2009;101:430–437.
 54. Narres M, Claessen H, Droste S, Kvitkina T, Koch M, Kuss O, Icks A. The incidence of end-stage renal disease in the diabetic (compared to the non-diabetic) population: a systematic review. *PLoS One*. 2016;11:e0147329. doi: 10.1371/journal.pone.0147329.
 55. Lopes AA. End-stage renal disease due to diabetes in racial/ethnic minorities and disadvantaged populations. *Ethn Dis*. 2009;19(suppl 1):S1–S47.
 56. Lefebvre KM, Lavery LA. Disparities in amputations in minorities. *Clin Orthop Relat Res*. 2011;469:1941–1950. doi: 10.1007/s11999-011-1842-x.
 57. Taylor HA Jr, Akylbekova EL, Garrison RJ, Sarpong D, Joe J, Walker E, Wyatt SB, Steffes MW. Dyslipidemia and the treatment of lipid disorders in African Americans. *Am J Med*. 2009;122:454–463. doi: 10.1016/j.amjmed.2008.09.049.
 58. Zweifler RM, McClure LA, Howard VJ, Cushman M, Hovater MK, Safford MM, Howard G, Goff DC Jr. Racial and geographic differences in prevalence, awareness, treatment and control of dyslipidemia: the Reasons for

- Geographic and Racial Differences in Stroke (REGARDS) study. *Neuroepidemiology*. 2011;37:39–44. doi: 10.1159/000328258.
59. Goff DC Jr, Bertoni AG, Kramer H, Bonds D, Blumenthal RS, Tsai MY, Psaty BM. Dyslipidemia prevalence, treatment, and control in the Multi-Ethnic Study of Atherosclerosis (MESA): gender, ethnicity, and coronary artery calcium. *Circulation*. 2006;113:647–656. doi: 10.1161/CIRCULATIONAHA.105.552737.
 60. Stone NJ, Robinson J, Lichtenstein AH, Bairey Merz CN, Blum CB, Eckel RH, Goldberger AC, Gordon D, Levy D, Lloyd-Jones DM, McBride P, Schwartz JS, Shero ST, Smith SC Jr, Watson K, Wilson PW. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines [published corrections appear in *Circulation*. 2014;129[suppl 2]:S46–S48 and *Circulation*. 2015;132:e396]. *Circulation*. 2014;129(suppl 2):S1–S45. doi: 10.1161/01.cir.0000437738.63853.7a.
 61. Ogden CL, Carroll MD, Lawman HG, Fryar CD, Kruszon-Moran D, Kit BK, Flegal KM. Trends in obesity prevalence among children and adolescents in the United States, 1988–1994 through 2013–2014. *JAMA*. 2016;315:2292–2299. doi: 10.1001/jama.2016.6361.
 62. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA*. 2014;311:806–814. doi: 10.1001/jama.2014.732.
 63. Calle EE, Thun MJ, Petrelli JM, Rodriguez C, Heath CW Jr. Body-mass index and mortality in a prospective cohort of U.S. adults. *N Engl J Med*. 1999;341:1097–1105. doi: 10.1056/NEJM199910073411501.
 64. Standl E, Erbach M, Schnell O. Defending the con side: obesity paradox does not exist. *Diabetes Care*. 2013;36(suppl 2):S282–S286. doi: 10.2337/dcS13-2040.
 65. McDowell MA, Fryar CD, Ogden CL, Flegal KM. Anthropometric reference data for children and adults: United States, 2003–2006. *Natl Health Stat Report*. 2008;22:1–48.
 66. Carroll JF, Chiapa AL, Rodriguez M, Phelps DR, Cardarelli KM, Vishwanatha JK, Bae S, Cardarelli R. Visceral fat, waist circumference, and BMI: impact of race/ethnicity. *Obesity (Silver Spring)*. 2008;16:600–607. doi: 10.1038/oby.2007.92.
 67. Deleted in proof.
 68. Hill JO, Sidney S, Lewis CE, Tolan K, Scherzinger AL, Stamm ER. Racial differences in amounts of visceral adipose tissue in young adults: the CARDIA (Coronary Artery Risk Development in Young Adults) study. *Am J Clin Nutr*. 1999;69:381–387.
 69. Després JP, Lemieux I. Abdominal obesity and metabolic syndrome. *Nature*. 2006;444:881–887. doi: 10.1038/nature05488.
 70. Katzmarzyk PT, Bray GA, Greenway FL, Johnson WD, Newton RL Jr, Ravussin E, Ryan DH, Smith SR, Bouchard C. Racial differences in abdominal depot-specific adiposity in white and African American adults. *Am J Clin Nutr*. 2010;91:7–15. doi: 10.3945/ajcn.2009.28136.
 71. Langellier BA, Glik D, Ortega AN, Prelep ML. Trends in racial/ethnic disparities in overweight self-perception among US adults, 1988–1994 and 1999–2008. *Public Health Nutr*. 2015;18:2115–2125. doi: 10.1017/S1368980014002560.
 72. Bennett GG, Wolin KY. Satisfied or unaware? Racial differences in perceived weight status. *Int J Behav Nutr Phys Act*. 2006;3:40. doi: 10.1186/1479-5868-3-40.
 73. Duncan DT, Wolin KY, Scharoun-Lee M, Ding EL, Warner ET, Bennett GG. Does perception equal reality? Weight misperception in relation to weight-related attitudes and behaviors among overweight and obese US adults. *Int J Behav Nutr Phys Act*. 2011;8:20. doi: 10.1186/1479-5868-8-20.
 74. Lynch E, Liu K, Wei GS, Spring B, Kiefe C, Greenland P. The relation between body size perception and change in body mass index over 13 years: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Am J Epidemiol*. 2009;169:857–866. doi: 10.1093/aje/kwn412.
 75. Williamson DF, Serdula MK, Anda RF, Levy A, Byers T. Weight loss attempts in adults: goals, duration, and rate of weight loss. *Am J Public Health*. 1992;82:1251–1257.
 76. Wolf PA, D'Agostino RB, Belanger AJ, Kannel WB. Probability of stroke: a risk profile from the Framingham Study. *Stroke*. 1991;22:312–318.
 77. Manolio TA, Kronmal RA, Burke GL, O'Leary DH, Price TR. Short-term predictors of incident stroke in older adults: the Cardiovascular Health Study. *Stroke*. 1996;27:1479–1486.
 78. Soliman EZ, Safford MM, Muntner P, Khodneva Y, Dawood FZ, Zakai NA, Thacker EL, Judd S, Howard VJ, Howard G, Herrington DM, Cushman M. Atrial fibrillation and the risk of myocardial infarction [published correction appears in *JAMA Intern Med*. 2014;174:308]. *JAMA Intern Med*. 2014;174:107–114. doi: 10.1001/jamainternmed.2013.11912.
 79. Soliman EZ, Lopez F, O'Neal WT, Chen LY, Bengtson L, Zhang ZM, Loehr L, Cushman M, Alonso A. Atrial fibrillation and risk of ST-segment-elevation versus non-ST-segment-elevation myocardial infarction: the Atherosclerosis Risk in Communities (ARIC) Study. *Circulation*. 2015;131:1843–1850. doi: 10.1161/CIRCULATIONAHA.114.014145.
 80. Violi F, Soliman EZ, Pignatelli P, Pastori D. Atrial fibrillation and myocardial infarction: a systematic review and appraisal of pathophysiologic mechanisms. *J Am Heart Assoc*. 2016;5: e003347. doi: 10.1161/JAHA.116.003347.
 81. Soliman EZ, Alonso A, Goff DC Jr. Atrial fibrillation and ethnicity: the known, the unknown and the paradox. *Future Cardiol*. 2009;5:547–556. doi: 10.2217/fca.09.49.
 82. Alonso A, Agarwal SK, Soliman EZ, Ambrose M, Chamberlain AM, Prineas RJ, Folsom AR. Incidence of atrial fibrillation in whites and African-Americans: the Atherosclerosis Risk in Communities (ARIC) study. *Am Heart J*. 2009;158:111–117. doi: 10.1016/j.ahj.2009.05.010.
 83. Psaty BM, Manolio TA, Kuller LH, Kronmal RA, Cushman M, Fried LP, White R, Furberg CD, Rautaharju PM. Incidence of and risk factors for atrial fibrillation in older adults. *Circulation*. 1997;96:2455–2461.
 84. Meschia JF, Merrill P, Soliman EZ, Howard VJ, Barrett KM, Zakai NA, Kleindorfer D, Safford M, Howard G. Racial disparities in awareness and treatment of atrial fibrillation: the REasons for Geographic and Racial Differences in Stroke (REGARDS) study. *Stroke*. 2010;41:581–587. doi: 10.1161/STROKEAHA.109.573907.
 85. Krebs-Smith SM, Reedy J, Bosire C. Healthfulness of the U.S. food supply: little improvement despite decades of dietary guidance. *Am J Prev Med*. 2010;38:472–477. doi: 10.1016/j.amepre.2010.01.016.
 86. Jefferson WK, Zunker C, Feucht JC, Fitzpatrick SL, Greene LF, Shewchuk RM, Baskin ML, Walton NW, Phillips B, Ard JD. Use of the Nominal Group Technique (NGT) to understand the perceptions of the healthiness of foods associated with African Americans. *Eval Program Plann*. 2010;33:343–348. doi: 10.1016/j.evalprogplan.2009.11.002.
 87. Judd SE, Gutiérrez OM, Newby PK, Howard G, Howard VJ, Locher JL, Kissela BM, Shikany JM. Dietary patterns are associated with incident stroke and contribute to excess risk of stroke in black Americans. *Stroke*. 2013;44:3305–3311. doi: 10.1161/STROKEAHA.113.002636.
 88. Shikany JM, Safford MM, Newby PK, Durant RW, Brown TM, Judd SE. Southern dietary pattern is associated with hazard of acute coronary heart disease in the Reasons for Geographic and Racial Differences in Stroke (REGARDS) Study. *Circulation*. 2015;132:804–814. doi: 10.1161/CIRCULATIONAHA.114.014421.
 89. Cofta-Woerpel L, Randhawa V, McFadden HG, Fought A, Bullard E, Spring B. ACCISS study rationale and design: Activating Collaborative Cancer Information Service Support for cervical cancer screening. *BMC Public Health*. 2009;9:444. doi: 10.1186/1471-2458-9-444.
 90. Troiano RP, Berrigan D, Dodd KW, Mâsse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008;40:181–188. doi: 10.1249/mss.0b013e31815a51b3.
 91. Sidney S, Sternfeld B, Haskell WL, Jacobs DR Jr, Chesney MA, Hulley SB. Television viewing and cardiovascular risk factors in young adults: the CARDIA study. *Ann Epidemiol*. 1996;6:154–159.
 92. Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM, Pate RR, Troiano RP. Amount of time spent in sedentary behaviors in the United States, 2003–2004. *Am J Epidemiol*. 2008;167:875–881. doi: 10.1093/aje/kwm390.
 93. Hooker SP, Hutto B, Zhu W, Blair SN, Colabianchi N, Vena JE, Rhodes D, Howard VJ. Accelerometer measured sedentary behavior and physical activity in white and black adults: the REGARDS study. *J Sci Med Sport*. 2016;19:336–341. doi: 10.1016/j.jsms.2015.04.006.
 94. Airhihenbuwa CO, Kumanyika S, Agurs LC, Lowe A. Perceptions and beliefs about exercise, rest, and health among African-Americans. *Am J Health Promot*. 1995;9:426–429.
 95. Hooker SP, Wilson DK, Griffin SF, Ainsworth BE. Perceptions of environmental supports for physical activity in African American and white adults in a rural county in South Carolina. *Prev Chronic Dis*. 2005;2:A11.
 96. Resnicow K, Jackson A, Braithwaite R, Dilorio C, Blisset D, Rahotep S, Periasamy S. Healthy Body/Healthy Spirit: a church-based nutrition and physical activity intervention. *Health Educ Res*. 2002;17:562–573.
 97. Barnes AS, Goodrick GK, Pavlik V, Markesino J, Laws DY, Taylor WC. Weight loss maintenance in African-American women: focus group results and questionnaire development. *J Gen Intern Med*. 2007;22:915–922. doi: 10.1007/s11606-007-0195-3.
 98. Baskin ML, Ahluwalia HK, Resnicow K. Obesity intervention among African-American children and adolescents. *Pediatr Clin North Am*. 2001;48:1027–1039.

99. Joseph RP, Ainsworth BE, Keller C, Dodgson JE. Barriers to physical activity among African American women: an integrative review of the literature. *Women Health*. 2015;55:679–699. doi: 10.1080/03630242.2015.1039184.
100. Sanderson BK, Foushee HR, Bittner V, Cornell CE, Stalker V, Shelton S, Pulley L. Personal, social, and physical environmental correlates of physical activity in rural African-American women in Alabama. *Am J Prev Med*. 2003;25(suppl 1):30–37.
101. Rohm Young D, Voorhees CC. Personal, social, and environmental correlates of physical activity in urban African-American women. *Am J Prev Med*. 2003;25(suppl 1):38–44.
102. Huxley RR, Yatsuya H, Lutsey PL, Woodward M, Alonso A, Folsom AR. Impact of age at smoking initiation, dosage, and time since quitting on cardiovascular disease in African Americans and whites: the Atherosclerosis Risk in Communities Study. *Am J Epidemiol*. 2012;175:816–826. doi: 10.1093/aje/kwr391.
103. Garrett BE, Dube SR, Winder C, Caraballo RS, Centers for Disease Control and Prevention (CDC). Cigarette smoking—United States, 2006–2008 and 2009–2010. *MMWR Suppl*. 2013;62(suppl 3):81–84.
104. Eaton DK, Kann L, Kinchen S, Shanklin S, Flint KH, Hawkins J, Harris WA, Lowry R, McManus T, Chyen D, Whittle L, Lim C, Wechsler H; Centers for Disease Control and Prevention (CDC). Youth risk behavior surveillance—United States, 2011. *MMWR Surveill Summ*. 2012;61:1–162.
105. Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, Luepker R, Mittleman M, Samet J, Smith SC Jr, Tager I. Air pollution and cardiovascular disease: a statement for healthcare professionals from the Expert Panel on Population and Prevention Science of the American Heart Association. *Circulation*. 2004;109:2655–2671. doi: 10.1161/01.CIR.0000128587.30041.C8.
106. Iribarren C, Friedman GD, Klatsky AL, Eisner MD. Exposure to environmental tobacco smoke: association with personal characteristics and self reported health conditions. *J Epidemiol Community Health*. 2001;55:721–728.
107. Mannino DM, Siegel M, Rose D, Nkuchia J, Etzel R. Environmental tobacco smoke exposure in the home and worksite and health effects in adults: results from the 1991 National Health Interview Survey. *Tob Control*. 1997;6:296–305.
108. Steenland K, Sieber K, Etzel RA, Pechacek T, Maurer K. Exposure to environmental tobacco smoke and risk factors for heart disease among never smokers in the Third National Health and Nutrition Examination Survey. *Am J Epidemiol*. 1998;147:932–939.
109. Kabbani N. Not so cool? Menthol's discovered actions on the nicotinic receptor and its implications for nicotine addiction. *Front Pharmacol*. 2013;4:95. doi: 10.3389/fphar.2013.00095.
110. Richardson A, Ganz O, Pearson J, Celis N, Vallone D, Villanti AC. How the industry is marketing menthol cigarettes: the audience, the message and the medium. *Tob Control*. 2015;24:594–600. doi: 10.1136/tobaccocontrol-2014-051657.
111. Kasza KA, Hyland AJ, Bansal-Travers M, Vogl LM, Chen J, Evans SE, Fong GT, Cummings KM, O'Connor RJ. Switching between menthol and non-menthol cigarettes: findings from the U.S. Cohort of the International Tobacco Control Four Country Survey. *Nicotine Tob Res*. 2014;16:1255–1265. doi: 10.1093/ntr/ntu098.
112. Delnevo CD, Gundersen DA, Hrywna M, Echeverria SE, Steinberg MB. Smoking-cessation prevalence among U.S. smokers of menthol versus non-menthol cigarettes. *Am J Prev Med*. 2011;41:357–365. doi: 10.1016/j.amepre.2011.06.039.
113. St-Onge MP, Grandner MA, Brown D, Conroy MB, Jean-Louis G, Coons M, Bhatt DL; on behalf of the American Heart Association Obesity, Behavior Change, Diabetes, and Nutrition Committees of the Council on Lifestyle and Cardiometabolic Health; Council on Cardiovascular Disease in the Young; Council on Clinical Cardiology; Stroke Council. Sleep duration and quality: impact on lifestyle behaviors and cardiometabolic health: a scientific statement from the American Heart Association. *Circulation*. 2016;134:e367–e386. doi: 10.1161/CIR.0000000000000444.
114. Rangaraj VR, Knutson KL. Association between sleep deficiency and cardiometabolic disease: implications for health disparities. *Sleep Med*. 2016;18:19–35. doi: 10.1016/j.sleep.2015.02.535.
115. Ruiter ME, DeCoster J, Jacobs L, Lichstein KL. Sleep disorders in African Americans and Caucasian Americans: a meta-analysis. *Behav Sleep Med*. 2010;8:246–259. doi: 10.1080/15402002.2010.509251.
116. Patyar S, Patyar RR. Correlation between sleep duration and risk of stroke. *J Stroke Cerebrovasc Dis*. 2015;24:905–911. doi: 10.1016/j.jstrokecerebrovasdis.2014.12.038.
117. Knutson KL. Sleep duration and cardiometabolic risk: a review of the epidemiologic evidence. *Best Pract Res Clin Endocrinol Metab*. 2010;24:731–743. doi: 10.1016/j.beem.2010.07.001.
118. Liu TZ, Xu C, Rota M, Cai H, Zhang C, Shi MJ, Yuan RX, Weng H, Meng XY, Kwong JS, Sun X. Sleep duration and risk of all-cause mortality: a flexible, non-linear, meta-regression of 40 prospective cohort studies. *Sleep Med Rev*. 2017;32:28–36. doi: 10.1016/j.smrv.2016.02.005.
119. Covassin N, Singh P. Sleep duration and cardiovascular disease risk: epidemiologic and experimental evidence. *Sleep Med Clin*. 2016;11:81–89. doi: 10.1016/j.jsmc.2015.10.007.
120. Hale L, Do DP. Racial differences in self-reports of sleep duration in a population-based study. *Sleep*. 2007;30:1096–1103.
121. Grandner MA, Petrov ME, Rattanaumpawan P, Jackson N, Platt A, Patel NP. Sleep symptoms, race/ethnicity, and socioeconomic position. *J Clin Sleep Med*. 2013;9:897–905; 905A–905D. doi: 10.5664/jcs.m.2990.
122. Carnethon MR, De Chavez PJ, Zee PC, Kim KY, Liu K, Goldberger JJ, Ng J, Knutson KL. Disparities in sleep characteristics by race/ethnicity in a population-based sample: Chicago Area Sleep Study. *Sleep Med*. 2016;18:50–55. doi: 10.1016/j.sleep.2015.07.005.
123. Rasmussen-Torvik L, De Chavez PJ, Kershaw K, Knutson KL, Kim KA, Zee PC, Carnethon MR. The mediation of racial differences in hypertension by sleep characteristics: Chicago Area Sleep Study. *Am J Hypertens*. 2016;29:1353–1357. doi: 10.1093/ajh/hpw093.
124. Johnson DA, Lisabeth L, Hickson D, Johnson-Lawrence V, Samdarshi T, Taylor H, Diez Roux AV. The social patterning of sleep in African Americans: associations of socioeconomic position and neighborhood characteristics with sleep in the Jackson Heart Study. *Sleep*. 2016;39:1749–1759. doi: 10.5665/sleep.6106.
125. Johnson DA, Lisabeth L, Lewis TT, Sims M, Hickson DA, Samdarshi T, Taylor H, Diez Roux AV. The contribution of psychosocial stressors to sleep among African Americans in the Jackson Heart Study. *Sleep*. 2016;39:1411–1419. doi: 10.5665/sleep.5974.
126. Akinseye OA, Ojike NI, Akinseye LI, Dhandapani PS, Pandi-Perumal SR. Association of sleep duration with stroke in diabetic patients: analysis of the National Health Interview Survey. *J Stroke Cerebrovasc Dis*. 2016;25:650–655. doi: 10.1016/j.jstrokecerebrovasdis.2015.11.023.
127. Pearson TA, Palaniappan LP, Artinian NT, Carnethon MR, Criqui MH, Daniels SR, Fonarow GC, Fortmann SP, Franklin BA, Galloway JM, Goff DC Jr, Heath GW, Frank AT, Kris-Etherton PM, Labarthe DR, Murabito JM, Sacco RL, Sasson C, Turner MB; on behalf of the American Heart Association Council on Epidemiology and Prevention. American Heart Association guide for improving cardiovascular health at the community level, 2013 update: a scientific statement for public health practitioners, healthcare providers, and health policy makers. *Circulation*. 2013;127:1730–1753. doi: 10.1161/CIR.0b013e31828f8a94.
128. Kidney Disease Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl*. 2013;3:1–150.
129. Coresh J, Selvin E, Stevens LA, Manzi J, Kusek JW, Eggers P, Van Lente F, Levey AS. Prevalence of chronic kidney disease in the United States. *JAMA*. 2007;298:2038–2047. doi: 10.1001/jama.298.17.2038.
130. Grams ME, Juraschek SP, Selvin E, Foster MC, Inker LA, Eckfeldt JH, Levey AS, Coresh J. Trends in the prevalence of reduced GFR in the United States: a comparison of creatinine- and cystatin C-based estimates. *Am J Kidney Dis*. 2013;62:253–260. doi: 10.1053/j.ajkd.2013.03.013.
131. Tarver-Carr ME, Powe NR, Eberhardt MS, LaVeist TA, Kington RS, Coresh J, Brancati FL. Excess risk of chronic kidney disease among African-American versus white subjects in the United States: a population-based study of potential explanatory factors. *J Am Soc Nephrol*. 2002;13:2363–2370.
132. Muntner P, Newsome B, Kramer H, Peralta CA, Kim Y, Jacobs DR Jr, Kiefe CI, Lewis CE. Racial differences in the incidence of chronic kidney disease. *Clin J Am Soc Nephrol*. 2012;7:101–107. doi: 10.2215/CJN.06450611.
133. Crews DC, Liu Y, Boulware LE. Disparities in the burden, outcomes, and care of chronic kidney disease. *Curr Opin Nephrol Hypertens*. 2014;23:298–305. doi: 10.1097/01.mnh.0000444822.25991.16.
134. Freedman BI, Skorecki K. Gene-gene and gene-environment interactions in apolipoprotein L1 gene-associated nephropathy. *Clin J Am Soc Nephrol*. 2014;9:2006–2013. doi: 10.2215/CJN.01330214.
135. Naik RP, Derebail VK, Grams ME, Franceschini N, Auer PL, Peloso GM, Young BA, Lettre G, Peralta CA, Katz R, Hyacinth H, Quarells RC, Grove ML, Bick AG, Fontanillas P, Rich SS, Smith JD, Boerwinkle E, Rosamond WD, Ito K, Lanzkron S, Coresh J, Correa A, Sarto GE, Key NS, Jacobs DR, Kathiresan S, Bibbins-Domingo K, Kshirsagar AV, Wilson JG, Reiner AP.

- Association of sickle cell trait with chronic kidney disease and albuminuria in African Americans. *JAMA*. 2014;312:2115–2125. doi: 10.1001/jama.2014.15063.
136. Weiner DE, Tighiouart H, Amin MG, Stark PC, MacLeod B, Griffith JL, Salem DN, Levey AS, Sarnak MJ. Chronic kidney disease as a risk factor for cardiovascular disease and all-cause mortality: a pooled analysis of community-based studies. *J Am Soc Nephrol*. 2004;15:1307–1315.
 137. Gutiérrez OM, Khodneva YA, Muntner P, Rizk DV, McClellan WM, Cushman M, Warnock DG, Safford MM; REGARDS Investigators. Association between urinary albumin excretion and coronary heart disease in black vs white adults. *JAMA*. 2013;310:706–714. doi: 10.1001/jama.2013.8777.
 138. Gutiérrez OM, Judd SE, Muntner P, Rizk DV, McClellan WM, Safford MM, Cushman M, Kissela BM, Howard VJ, Warnock DG. Racial differences in albuminuria, kidney function, and risk of stroke. *Neurology*. 2012;79:1686–1692. doi: 10.1212/WNL.0b013e31826e9af8.
 139. Choi AI, Rodriguez RA, Bacchetti P, Bertenthal D, Hernandez GT, O'Hare AM. White/black racial differences in risk of end-stage renal disease and death. *Am J Med*. 2009;122:672–678. doi: 10.1016/j.amjmed.2008.11.021.
 140. US Renal Data System. USRDS 2010 annual data report: atlas of end-stage renal disease in the United States. 2010. <https://www.usrds.org/atlas10.aspx>. Accessed July 30, 2015.
 141. Newsome BB, McClellan WM, Coffey CS, Allison JJ, Kiefe CI, Warnock DG. Survival advantage of black patients with kidney disease after acute myocardial infarction. *Clin J Am Soc Nephrol*. 2006;1:993–999. doi: 10.2215/CJN.01251005.
 142. Crews DC, Sozio SM, Liu Y, Coresh J, Powe NR. Inflammation and the paradox of racial differences in dialysis survival. *J Am Soc Nephrol*. 2011;22:2279–2286. doi: 10.1681/ASN.2011030305.
 143. Jacobs AS, Ayinde HO, Lee DL. Inflammatory biomarkers and cardiovascular complications in sickle cell disease: a review. *Curr Cardiovasc Risk Rep*. 2013;7:368–377.
 144. Sandhu MK, Cohen A. Aging in sickle cell disease: co-morbidities and new issues in management. *Hemoglobin*. 2015;39:221–224. doi: 10.1009/03630269.2015.1040493.
 145. Mancini EA, Culbertson DE, Yang YM, Gardner TM, Powell R, Haynes J Jr, Shah AK, Mankad VN; Investigators of the Cooperative Study of Sickle Cell Disease. Causes of death in sickle cell disease: an autopsy study. *Br J Haematol*. 2003;123:359–365.
 146. Liem RI, Young LT, Thompson AA. Prolonged QTc interval in children and young adults with sickle cell disease at steady state. *Pediatr Blood Cancer*. 2009;52:842–846. doi: 10.1002/pbc.21973.
 147. Sangkatumvong S, Coates TD, Khoo MC. Abnormal autonomic cardiac response to transient hypoxia in sickle cell anemia. *Physiol Meas*. 2008;29:655–668. doi: 10.1088/0967-3334/29/5/010.
 148. Liem RI, Chan C, Vu TT, Fornage M, Thompson AA, Liu K, Carnethon MR. Association among sickle cell trait, fitness, and cardiovascular disease risk factors in Cardia. *Blood*. 2017;129:723–728. doi: 10.1182/blood-2016-07-727719.
 149. Caughey MC, Loehr LR, Key NS, Derebail VK, Gottesman RF, Kshirsagar AV, Grove ML, Heiss G. Sickle cell trait and incident ischemic stroke in the Atherosclerosis Risk in Communities study. *Stroke*. 2014;45:2863–2867. doi: 10.1161/STROKEAHA.114.006110.
 150. Harris KM, Haas TS, Eichner ER, Maron BJ. Sickle cell trait associated with sudden death in competitive athletes. *Am J Cardiol*. 2012;110:1185–1188. doi: 10.1016/j.amjcard.2012.06.004.
 151. Hyacinth HI, Cara CL, Seals SR, Irvin MR, Naik RP, Caughey M, Winkler C, Franceschini N, Burke GL, Zakai NA, Kopp JB, Judd SE, Adams RJ, Gee BE, Longstreth W, Egede LE, Lackland DT, Greenberg CS, Herman T, Manson JE, Key NS, Derebail VK, Kshirsagar AV, Folsom AR, Konety SH, Howard VJ, Allison M, Wilson JG, Correa A, Zhi D, Arnett D, Howard G, Cushman M, Reiner A, Safford MM. Association of sickle cell trait with risk of coronary heart disease in African Americans. *Blood*. 2016;128:11.
 152. Currier JS, Lundgren JD, Carr A, Klein D, Sabin CA, Sax PE, Schouten JT, Smieja M; Working Group 2. Epidemiological evidence for cardiovascular disease in HIV-infected patients and relationship to highly active antiretroviral therapy [published correction appears in *Circulation*. 2008;118:e108]. *Circulation*. 2008;118:e29–e35. doi: 10.1161/CIRCULATIONAHA.107.189624.
 153. Harris KM, Haas TS, Eichner ER, Maron BJ. Sickle cell trait associated with sudden death in competitive athletes. *Am J Cardiol*. 2012;110:1185–1188.
 154. Oramasionwu CU, Hunter JM, Brown CM, Morse GD, Lawson KA, Koeller JM, Frei CR. Cardiovascular disease in blacks with HIV/AIDS in the United States: a systematic review of the literature. *Open AIDS J*. 2012;6:29–35. doi: 10.2174/1874613601206010029.
 155. Oramasionwu CU, Morse GD, Lawson KA, Brown CM, Koeller JM, Frei CR. Hospitalizations for cardiovascular disease in African Americans and whites with HIV/AIDS. *Popul Health Manag*. 2013;16:201–207. doi: 10.1089/pop.2012.0043.
 156. Miller PE, Budoff M, Zikusoka M, Li X, Palella F Jr, Kingsley LA, Witt MD, Sharrett AR, Jacobson LP, Post WS. Comparison of racial differences in plaque composition and stenosis between HIV-positive and HIV-negative men from the Multicenter AIDS Cohort Study. *Am J Cardiol*. 2014;114:369–375. doi: 10.1016/j.amjcard.2014.04.049.
 157. Asztalos BF, Matera R, Horvath KV, Horan M, Tani M, Polak JF, Skinner S, Wanke CA. Cardiovascular disease-risk markers in HIV patients. *J AIDS Clin Res*. 2014;5:317.
 158. Botha S, Fourie CM, van Rooyen JM, Kruger A, Schutte AE. Cardiometabolic changes in treated versus never treated HIV-infected black South Africans: the PURE study. *Heart Lung Circ*. 2014;23:119–126. doi: 10.1016/j.hlc.2013.07.019.
 159. Poulter N. Coronary heart disease is a multifactorial disease. *Am J Hypertens*. 1999;12(pt 2):925–955.
 160. Ellis J, Lange EM, Li J, Dupuis J, Baumert J, Walston JD, Keating BJ, Durda P, Fox ER, Palmer CD, Meng YA, Young T, Farlow DN, Schnabel RB, Marzi CS, Larkin E, Martin LW, Bis JC, Auer P, Ramachandran VS, Gabriel SB, Willis MS, Pankow JS, Papanicolaou GJ, Rotter JJ, Ballantyne CM, Gross MD, Lettre G, Wilson JG, Peters U, Koenig W, Tracy RP, Redline S, Reiner AP, Benjamin EJ, Lange LA. Large multiethnic Candidate Gene Study for C-reactive protein levels: identification of a novel association at CD36 in African Americans. *Hum Genet*. 2014;133:985–995. doi: 10.1007/s00439-014-1439-z.
 161. Emerging Risk Factors Collaboration, Kaptoge S, Di Angelantonio E, Pennells L, Wood AM, White IR, Gao P, Walker M, Thompson A, Sarwar N, Caslake M, Butterworth AS, Amouyel P, Assmann G, Bakker SJ, Barr EL, Barrett-Connor E, Benjamin EJ, Björkelund C, Brenner H, Brunner E, Clarke R, Cooper JA, Cremer P, Cushman M, Dagenais GR, D'Agostino RB Sr, Dankner R, Davey-Smith G, Deeg D, Dekker JM, Engstrom G, Folsom AR, Fowkes FG, Gallacher J, Gaziano JM, Giampaoli S, Gillum RF, Hofman A, Howard BV, Ingelsson E, Iso H, Jorgensen T, Kiechl S, Kitamura A, Kiyohara Y, Koenig W, Kromhout D, Kuller LH, Lawlor DA, Meade TW, Nissinen A, Nordestgaard BG, Onat A, Panagiotakos DB, Psaty BM, Rodriguez B, Rosengren A, Salomaa V, Kauhanen J, Salonen JT, Shaffer JA, Shea S, Ford I, Stehouwer CD, Strandberg TE, Tipping RW, Toesetto A, Wassertheil-Smolter S, Wennberg P, Westendorp RG, Whincup PH, Wilhelmsen L, Woodward M, Lowe GD, Wareham NJ, Khaw KT, Sattar N, Packard CJ, Gudnason V, Ridker PM, Pepys MB, Thompson SG, Danesh J. C-reactive protein, fibrinogen, and cardiovascular disease prediction. *N Engl J Med*. 2012;367:1310–1320. doi: 10.1056/NEJMoa1107477.
 162. Reiner AP, Beleza S, Franceschini N, Auer PL, Robinson JG, Kooperberg C, Peters U, Tang H. Genome-wide association and population genetic analysis of C-reactive protein in African American and Hispanic American women. *Am J Hum Genet*. 2012;91:502–512. doi: 10.1016/j.ajhg.2012.07.023.
 163. Schick UM, Auer PL, Bis JC, Lin H, Wei P, Pankratz N, Lange LA, Brody J, Stitzel NO, Kim DS, Carlson CS, Fornage M, Haessler J, Hsu L, Jackson RD, Kooperberg C, Leal SM, Psaty BM, Boerwinkle E, Tracy R, Ardisson D, Shah S, Willer C, Loos R, Melander O, McPherson R, Hovingh K, Reilly M, Watkins H, Girelli D, Fontanillas P, Chasman DI, Gabriel SB, Gibbs R, Nickerson DA, Kathiresan S, Peters U, Dupuis J, Wilson JG, Rich SS, Morrison AC, Benjamin EJ, Gross MD, Reiner AP; Cohorts for Heart and Aging Research in Genomic Epidemiology; National Heart, Lung, and Blood Institute GO Exome Sequencing Project. Association of exome sequences with plasma C-reactive protein levels in >9000 participants. *Hum Mol Genet*. 2015;24:559–571. doi: 10.1093/hmg/ddu450.
 164. Herrick S, Blanc-Brude O, Gray A, Laurent G. Fibrinogen. *Int J Biochem Cell Biol*. 1999;31:741–746.
 165. Fibrinogen Studies Collaboration, Danesh J, Lewington S, Thompson SG, Lowe GD, Collins R, Kostis JB, Wilson AC, Folsom AR, Wu K, Bendersly M, Goldbourt U, Willeit J, Kiechl S, Yarnell JW, Sweetnam PM, Elwood PC, Cushman M, Psaty BM, Tracy RP, Tybjaerg-Hansen A, Haverkate F, de Maat MP, Fowkes FG, Lee AJ, Smith FB, Salomaa V, Harald K, Rasi R, Vahtera E, Jousilahti P, Pekkanen J, D'Agostino R, Kannel WB, Wilson PW, Tofler G, Arocha-Pinango CL, Rodriguez-Larralde A, Nagy E, Mijares M, Espinosa R, Rodriguez-Roa E, Ryder E, Diez-Ewald MP, Campos G, Fernandez V, Torres E, Marchioli R, Valagussa F, Rosengren A, Wilhelmsen L, Lappas G, Eriksson H, Cremer P, Nagel D, Curb JD, Rodriguez B, Yano

- K, Salonen JT, Nyyssönen K, Tuomainen TP, Hedblad B, Lind P, Loewel H, Koenig W, Meade TW, Cooper JA, De Stavola B, Knottenbelt C, Miller GJ, Cooper JA, Bauer KA, Rosenberg RD, Sato S, Kitamura A, Naito Y, Palosuo T, Ducimetiere P, Amouyel P, Arveiler D, Evans AE, Ferrières J, Juhan-Vague I, Bingham A, Schulte H, Assmann G, Cantin B, Lamarche B, Despres JP, Dagenais GR, Tunstall-Pedoe H, Woodward M, Ben-Shlomo Y, Davey Smith G, Palmieri V, Yeh JL, Rudnicka A, Ridker P, Rodeghiero F, Tosto A, Shepherd J, Ford I, Robertson M, Brunner E, Shipley M, Feskens EJ, Kromhout D, Dickinson A, Ireland B, Juzwishin K, Kaptoge S, Lewington S, Memon A, Sarwar N, Walker M, Wheeler J, White I, Wood A. Plasma fibrinogen level and the risk of major cardiovascular diseases and nonvascular mortality: an individual participant meta-analysis [published correction appears in *JAMA*. 2005;294:2848]. *JAMA*. 2005;294:1799–1809.
166. van Holten TC, Waanders LF, de Groot PG, Vissers J, Hoefer IE, Pasterkamp G, Prins MW, Roest M. Circulating biomarkers for predicting cardiovascular disease risk: a systematic review and comprehensive overview of meta-analyses. *PLoS One*. 2013;8:e62080. doi: 10.1371/journal.pone.0062080.
 167. Rjosk-Dendorfer D, Gürtler VM, Sommer WH, Reiser M, Clevert DA. Value of high resolution compression elastography and color Doppler sonography in characterisation of breast lesions: comparison of different high-frequency transducers. *Clin Hemorheol Microcirc*. 2014;57:129–135. doi: 10.3233/CH-141824.
 168. Sabater-Lleal M, Huang J, Chasman D, Naitza S, Dehghan A, Johnson AD, Teumer A, Reiner AP, Folkersen L, Basu S, Rudnicka AR, Trompet S, Mälarstig A, Baumert J, Bis JC, Guo X, Hottenga JJ, Shin SY, Lopez LM, Lahti J, Tanaka T, Yanek LR, Oudot-Mellakh T, Wilson JF, Navarro P, Huffman JE, Zemunik T, Redline S, Mehra R, Pulanic D, Rudan I, Wright AF, Kolcic I, Polasek O, Wild SH, Campbell H, Curb JD, Wallace R, Liu S, Eaton CB, Becker DM, Becker LC, Bandinelli S, Räikkönen K, Widen E, Palotie A, Fornage M, Green D, Gross M, Davies G, Harris SE, Liewald DC, Starr JM, Williams FM, Grant PJ, Spector TD, Strawbridge RJ, Silveira A, Sennblad B, Rivadeneira F, Uitterlinden AG, Franco OH, Hofman A, van Dongen J, Willemssen G, Boomsma DI, Yao J, Swords Jenny N, Haritunians T, McKnight B, Lumley T, Taylor KD, Rotter JJ, Psaty BM, Peters A, Gieger C, Illig T, Grotevendt A, Homuth G, Völzke H, Kocher T, Goel A, Franzosi MG, Seedorf U, Clarke R, Steri M, Tarasov KV, Sanna S, Schlessinger D, Stott DJ, Sattar N, Buckley BM, Rumley A, Lowe GD, McArdle WL, Chen MH, Tofler GH, Song J, Boerwinkle E, Folsom AR, Rose LM, Franco-Cereceda A, Teichert M, Ikram MA, Mosley TH, Bevan S, Dichgans M, Rothwell PM, Sudlow CL, Hopewell JC, Chambers JC, Saleheen D, Kooner JS, Danesh J, Nelson CP, Erdmann J, Reilly MP, Kathiresan S, Schunkert H, Morange PE, Ferrucci L, Eriksson JG, Jacobs D, Deary IJ, Soranzo N, Witteman JC, de Geus EJ, Tracy RP, Hayward C, Koenig W, Cucca F, Jukema JW, Eriksson P, Seshadri S, Markus HS, Watkins H, Samani NJ; VTE Consortium; STROKE Consortium; Wellcome Trust Case Control Consortium 2 (WTCCC2); C4D Consortium; CARDIoGRAM Consortium; Wallaschofski H, Smith NL, Tregouet D, Ridker PM, Tang W, Strachan DP, Hamsten A, O'Donnell CJ. Multiethnic meta-analysis of genome-wide association studies in >100 000 subjects identifies 23 fibrinogen-associated loci but no strong evidence of a causal association between circulating fibrinogen and cardiovascular disease. *Circulation*. 2013;128:1310–1324. doi: 10.1161/CIRCULATIONAHA.113.002251.
 169. Folsom AR, Wu KK, Conlan MG, Finch A, Davis CE, Marcucci G, Sorlie PD, Szklo M. Distributions of hemostatic variables in blacks and whites: population reference values from the Atherosclerosis Risk in Communities (ARIC) Study. *Ethn Dis*. 1992;2:35–46.
 170. Vorster HH, Jerling JC, Steyn K, Badenhorst CJ, Slazus W, Venter CS, Jooste PL, Bourne LT. Plasma fibrinogen of black South Africans: the BRISK study. *Public Health Nutr*. 1998;1:169–176.
 171. Fibrinogen Studies Collaboration, Kaptoge S, White IR, Thompson SG, Wood AM, Lewington S, Lowe GD, Danesh J. Associations of plasma fibrinogen levels with established cardiovascular disease risk factors, inflammatory markers, and other characteristics: individual participant meta-analysis of 154,211 adults in 31 prospective studies: the Fibrinogen Studies Collaboration. *Am J Epidemiol*. 2007;166:867–879.
 172. Nienaber C, Pieters M, Kruger SH, Stonehouse W, Vorster HH. Overfatness, stunting and physical inactivity are determinants of plasminogen activator inhibitor-1 activity, fibrinogen and thrombin-antithrombin complex in African adolescents. *Blood Coagul Fibrinolysis*. 2008;19:361–368. doi: 10.1097/MB.C.0b013e328304b61a.
 173. Pieters M, Kotze RC, Jerling JC, Kruger A, Ariëns RA. Evidence that fibrinogen γ' regulates plasma clot structure and lysis and relationship to cardiovascular risk factors in black Africans. *Blood*. 2013;121:3254–3260. doi: 10.1182/blood-2012-12-471482.
 174. Dehghan A, Yang Q, Peters A, Basu S, Bis JC, Rudnicka AR, Kavousi M, Chen MH, Baumert J, Lowe GD, McKnight B, Tang W, de Maat M, Larson MG, Eyhermendi S, McArdle WL, Lumley T, Pankow JS, Hofman A, Massaro JM, Rivadeneira F, Kolz M, Taylor KD, van Duijn CM, Kathiresan S, Illig T, Aulchenko YS, Volcik KA, Johnson AD, Uitterlinden AG, Tofler GH, Gieger C, Wellcome Trust Case Control Consortium, Psaty BM, Couper DJ, Boerwinkle E, Koenig W, O'Donnell CJ, Witteman JC, Strachan DP, Smith NL, Folsom AR. Association of novel genetic loci with circulating fibrinogen levels: a genome-wide association study in 6 population-based cohorts. *Circ Cardiovasc Genet*. 2009;2:125–133. doi: 10.1161/CIRCGENETICS.108.825224.
 175. Francis CW, Marder VJ, Martin SE. Demonstration of a large molecular weight variant of the gamma chain of normal human plasma fibrinogen. *J Biol Chem*. 1980;255:5599–5604.
 176. Pieters M, de Maat MP, Jerling JC, Hoekstra T, Kruger A. Fibrinogen concentration and its role in CVD risk in black South Africans: effect of urbanisation. *Thromb Haemost*. 2011;106:448–456. doi: 10.1160/TH11-03-0192.
 177. Reiner AP, Carty CL, Carlson CS, Wan JY, Rieder MJ, Smith JD, Rice K, Fornage M, Jaquish CE, Williams OD, Tracy RP, Lewis CE, Siscovick DS, Boerwinkle E, Nickerson DA. Association between patterns of nucleotide variation across the three fibrinogen genes and plasma fibrinogen levels: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *J Thromb Haemost*. 2006;4:1279–1287. doi: 10.1111/j.1538-7836.2006.01907.x.
 178. Wassel CL, Lange LA, Keating BJ, Taylor KC, Johnson AD, Palmer C, Ho LA, Smith NL, Lange EM, Li Y, Yang Q, Delaney JA, Tang W, Tofler G, Redline S, Taylor HA Jr, Wilson JG, Tracy RP, Jacobs DR Jr, Folsom AR, Green D, O'Donnell CJ, Reiner AP. Association of genomic loci from a cardiovascular gene SNP array with fibrinogen levels in European Americans and African-Americans from six cohort studies: the Candidate Gene Association Resource (CARE). *Blood*. 2011;117:268–275. doi: 10.1182/blood-2010-06-289546.
 179. Jeff JM, Brown-Gentry K, Crawford DC. Replication and characterisation of genetic variants in the fibrinogen gene cluster with plasma fibrinogen levels and haematological traits in the Third National Health and Nutrition Examination Survey. *Thromb Haemost*. 2012;107:458–467. doi: 10.1160/TH11-07-0497.
 180. Kotzé RC, Nienaber-Rousseau C, De Lange Z, De Maat MP, Hoekstra T, Pieters M. Genetic polymorphisms influencing total and γ' fibrinogen levels and fibrin clot properties in Africans. *Br J Haematol*. 2015;168:102–112. doi: 10.1111/bjh.13104.
 181. Hunt SC, Hasstedt SJ, Kuida H, Stults BM, Hopkins PN, Williams RR. Genetic heritability and common environmental components of resting and stressed blood pressures, lipids, and body mass index in Utah pedigrees and twins. *Am J Epidemiol*. 1989;129:625–638.
 182. Williams PD, Puddey IB, Martin NG, Beilin LJ. Genetic and environmental covariance of serum cholesterol and blood pressure in female twins. *Atherosclerosis*. 1993;100:19–31.
 183. Somes GW, Harshfield GA, Alpert BS, Goble MM, Schieken RM. Genetic influences on ambulatory blood pressure patterns: the Medical College of Virginia Twin Study. *Am J Hypertens*. 1995;8(pt 1):474–478.
 184. Biron P, Mongeau JG, Bertrand D. Familial aggregation of blood pressure in 558 adopted children. *Can Med Assoc J*. 1976;115:773–774.
 185. Williams RR, Hunt SC, Hasstedt SJ, Hopkins PN, Wu LL, Berry TD, Stults BM, Barlow GK, Schumacher MC, Lifton RP, Lalouel JM. Are there interactions and relations between genetic and environmental factors predisposing to high blood pressure? *Hypertension*. 1991;18(suppl):129–137.
 186. Hunt SC, Williams RR, Barlow GK. A comparison of positive family history definitions for defining risk of future disease [published correction appears in *J Chronic Dis*. 1987;40:369]. *J Chronic Dis*. 1986;39:809–821.
 187. Levy D, Ehret GB, Rice K, Verwoert GC, Launer LJ, Dehghan A, Glazer NL, Morrison AC, Johnson AD, Aspelund T, Aulchenko Y, Lumley T, Köttgen A, Vasan RS, Rivadeneira F, Eiriksdottir G, Guo X, Arking DE, Mitchell JB, Mattace-Raso FU, Smith AV, Taylor K, Scharpf RB, Hwang SJ, Sijbrands EJ, Bis J, Harris TB, Ganesh SK, O'Donnell CJ, Hofman A, Rotter JJ, Coresh J, Benjamin EJ, Uitterlinden AG, Heiss G, Fox CS, Witteman JC, Boerwinkle E, Wang TJ, Gudnason V, Larson MG, Chakravarti A, Psaty BM, van Duijn CM. Genome-wide association study of blood pressure and hypertension. *Nat Genet*. 2009;41:677–687. doi: 10.1038/ng.384.
 188. Ganesh SK, Tragante V, Guo W, Guo Y, Lanktree MB, Smith EN, Johnson T, Castillo BA, Barnard J, Baumert J, Chang YP, Elbers CC, Farrall M,

- Fischer ME, Franceschini N, Gaunt TR, Gho JM, Gieger C, Gong Y, Isaacs A, Kleber ME, Mateo Leach I, McDonough CW, Meijis MF, Mellander O, Molony CM, Nolte IM, Padmanabhan S, Price TS, Rajagopalan R, Shaffer J, Shah S, Shen H, Soranzo N, van der Most PJ, Van Iperen EP, Van Setten J, Van Setten JA, Vonk JM, Zhang L, Beiltshees AL, Berenson GS, Bhatt DL, Boer JM, Boerwinkle E, Burkley B, Burt A, Chakravarti A, Chen W, Cooper-Dehoff RM, Curtis SP, Dreisbach A, Duggan D, Ehret GB, Fabsitz RR, Fornage M, Fox E, Furlong CE, Gansevoort RT, Hofker MH, Hovingh GK, Kirkland SA, Kottke-Marchant K, Kutlar A, Lacroix AZ, Langa TY, Li YR, Lin H, Liu K, Maiwald S, Malik R; CARDIOGRAM, METASTROKE, Murugesan G, Newton-Cheh C, O'Connell JR, Onland-Moret NC, Ouwehand WH, Palmas W, Penninx BW, Pepine CJ, Pettinger M, Polak JF, Ramachandran VS, Ranchalis J, Redline S, Ridker PM, Rose LM, Scharnag H, Schork NJ, Shimbo D, Shuldiner AR, Srinivasan SR, Stolk RP, Taylor HA, Thorand B, Trip MD, van Duijn CM, Verschuren WM, Wijmenga C, Winkelman BR, Wyatt S, Young JH, Boehm BO, Caulfield MJ, Chasman DI, Davidson KW, Doevendans PA, Fitzgerald GA, Gums JG, Hakonarson H, Hillege HL, Illig T, Jarvik GP, Johnson JA, Kastelein JJ, Koenig W; LifeLines Cohort Study, März W, Mitchell BD, Murray SS, Oldehinkel AJ, Rader DJ, Reilly MP, Reiner AP, Schadt EE, Silverstein RR, Snieder H, Stanton AV, Uitterlinden AG, van der Harst P, van der Schouw YT, Samani NJ, Johnson AD, Munroe PB, de Bakker PI, Zhu X, Levy D, Keating BJ, Asselbergs FW. Loci influencing blood pressure identified using a cardiovascular gene-centric array. *Hum Mol Genet*. 2013;22:1663–1678. doi: 10.1093/hmg/dd555.
189. Adeyemo A, Gerry N, Chen G, Herbert A, Doumatey A, Huang H, Zhou J, Lashley K, Chen Y, Christman M, Rotimi C. A genome-wide association study of hypertension and blood pressure in African Americans. *PLoS Genet*. 2009;5:e1000564. doi: 10.1371/journal.pgen.1000564.
190. Fox ER, Young JH, Li Y, Dreisbach AW, Keating BJ, Musani SK, Liu K, Morrison AC, Ganesh S, Kutlar A, Ramachandran VS, Polak JF, Fabsitz RR, Dries DL, Farlow DN, Redline S, Adeyemo A, Hirschhorn JN, Sun YV, Wyatt SB, Penman AD, Palmas W, Rotter JJ, Townsend RR, Doumatey AP, Tayo BO, Mosley TH Jr, Lyon HN, Kang SJ, Rotimi CN, Cooper RS, Franceschini N, Curb JD, Martin LW, Eaton CB, Kardia SL, Taylor HA, Caulfield MJ, Ehret GB, Johnson T; International Consortium for Blood Pressure Genome-wide Association Studies (ICBP-GWAS), Chakravarti A, Zhu X, Levy D. Association of genetic variation with systolic and diastolic blood pressure among African Americans: the Candidate Gene Association Resource study. *Hum Mol Genet*. 2011;20:2273–2284. doi: 10.1093/hmg/ddr092.
191. Zhu X, Young JH, Fox E, Keating BJ, Franceschini N, Kang S, Tayo B, Adeyemo A, Sun YV, Li Y, Morrison A, Newton-Cheh C, Liu K, Ganesh SK, Kutlar A, Vasan RS, Dreisbach A, Wyatt S, Polak J, Palmas W, Musani S, Taylor H, Fabsitz R, Townsend RR, Dries D, Glessner J, Chiang CW, Mosley T, Kardia S, Curb D, Hirschhorn JN, Rotimi C, Reiner A, Eaton C, Rotter JJ, Cooper RS, Redline S, Chakravarti A, Levy D. Combined admixture mapping and association analysis identifies a novel blood pressure genetic locus on 5p13: contributions from the CaRe consortium. *Hum Mol Genet*. 2011;20:2285–2295. doi: 10.1093/hmg/ddr113.
192. Kidambi S, Ghosh S, Kotchen JM, Grim CE, Krishnaswami S, Kaldunski ML, Cowley AW Jr, Patel SB, Kotchen TA. Non-replication study of a genome-wide association study for hypertension and blood pressure in African Americans. *BMC Med Genet*. 2012;13:27. doi: 10.1186/1471-2350-13-27.
193. Tran NT, Aslibekyan S, Tiwari HK, Zhi D, Sung YJ, Hunt SC, Rao DC, Broeckel U, Judd SE, Muntner P, Kent ST, Arnett DK, Irvin MR. PCSK9 variation and association with blood pressure in African Americans: preliminary findings from the HyperGEN and REGARDS studies. *Front Genet*. 2015;6:136. doi: 10.3389/fgene.2015.00136.
194. Liu C, Kraja AT, Smith JA, Brody JA, Franceschini N, Bis JC, Rice K, Morrison AC, Lu Y, Weiss S, Guo X, Palmas W, Martin LW, Chen YD, Surendran P, Drenos F, Cook JP, Auer PL, Chu AY, Giri A, Zhao W, Jakobsdottir J, Lin LA, Stafford JM, Amin N, Mei H, Yao J, Voorman A; CHD Exome+ Consortium; ExomeBP Consortium; GoT2DGenes Consortium; T2D-GENES Consortium; Larson MG, Grove ML, Smith AV, Hwang SJ, Chen H, Huan T, Kosova G, Stitzel NO, Kathiresan S, Samani N, Schunkert H, Deloukas P; Myocardial Infarction Genetics and CARDIoGRAM Exome Consortia, Li M, Fuchsberger C, Pattaro C, Gorski M; CKDGen Consortium, Kooperberg C, Papanicolaou GJ, Rossouw JE, Faul JD, Kardia SL, Bouchard C, Raffel LJ, Uitterlinden AG, Franco OH, Vasan RS, O'Donnell CJ, Taylor KD, Liu K, Bottinger EP, Gottesman O, Daw EW, Giulianini F, Ganesh S, Salfati E, Harris TB, Launer LJ, Dörr M, Felix SB, Rettig R, Völzke H, Kim E, Lee WJ, Lee IT, Sheu WH, Tsosie KS, Edwards DR, Liu Y, Correa A, Weir DR, Völker U, Ridker PM, Boerwinkle E, Gudnason V, Reiner AP, van Duijn CM, Borecki IB, Edwards TL, Chakravarti A, Rotter JJ, Psaty BM, Loos RJ, Fornage M, Ehret GB, Newton-Cheh C, Levy D, Chasman DI. Meta-analysis identifies common and rare variants influencing blood pressure and overlapping with metabolic trait loci. *Nat Genet*. 2016;48:1162–1170. doi: 10.1038/ng.3660.
195. Keebler ME, Sanders CL, Surti A, Guiducci C, Burt NP, Kathiresan S. Association of blood lipids with common DNA sequence variants at 19 genetic loci in the multiethnic United States National Health and Nutrition Examination Survey III. *Circ Cardiovasc Genet*. 2009;2:238–243. doi: 10.1161/CIRCGENETICS.108.829473.
196. Chang MH, Yesupriya A, Ned RM, Mueller PW, Dowling NF. Genetic variants associated with fasting blood lipids in the U.S. population: Third National Health and Nutrition Examination Survey. *BMC Med Genet*. 2010;11:62. doi: 10.1186/1471-2350-11-62.
197. Chang MH, Ned RM, Hong Y, Yesupriya A, Yang Q, Liu T, Janssens AC, Dowling NF. Racial/ethnic variation in the association of lipid-related genetic variants with blood lipids in the US adult population. *Circ Cardiovasc Genet*. 2011;4:523–533. doi: 10.1161/CIRCGENETICS.111.959577.
198. TG and HDL Working Group of the Exome Sequencing Project, National Heart, Lung, and Blood Institute, Crosby J, Peloso GM, Auer PL, Crosslin DR, Stitzel NO, Lange LA, Lu Y, Tang ZZ, Zhang H, Hindy G, Masca N, Stirrups K, Kanoni S, Do R, Jun G, Hu Y, Kang HM, Xue C, Goel A, Farrall M, Duga S, Merlini PA, Asselta R, Girelli D, Olivieri O, Martinelli N, Yin W, Reilly D, Speliotes E, Fox CS, Hveem K, Holmen OL, Nikpay M, Farlow DN, Assimes TL, Franceschini N, Robinson J, North KE, Martin LW, DePristo M, Gupta N, Escher SA, Jansson JH, Van Zuydam N, Palmer CN, Wareham N, Koch W, Meitinger T, Peters A, Lieb W, Erbel R, König IR, Kruppa J, Degenhardt F, Gottesman O, Bottinger EP, O'Donnell CJ, Psaty BM, Ballantyne CM, Abecasis G, Ordovas JM, Melander O, Watkins H, Orho-Melander M, Ardisson D, Loos RJ, McPherson R, Willer CJ, Erdmann J, Hall AS, Samani NJ, Deloukas P, Schunkert H, Wilson JG, Kooperberg C, Rich SS, Tracy RP, Lin DY, Altschuler D, Gabriel S, Nickerson DA, Jarvik GP, Cupples LA, Reiner AP, Boerwinkle E, Kathiresan S. Loss-of-function mutations in APOC3, triglycerides, and coronary disease. *N Engl J Med*. 2014;371:22–31.
199. Ito K, Bick AG, Flannick J, Friedman DJ, Genovese G, Parfenov MG, Depalma SR, Gupta N, Gabriel SB, Taylor HA Jr, Fox ER, Newton-Cheh C, Kathiresan S, Hirschhorn JN, Altschuler DM, Pollak MR, Wilson JG, Seidman JG, Seidman C. Increased burden of cardiovascular disease in carriers of APO11 genetic variants. *Circ Res*. 2014;114:845–850. doi: 10.1161/CIRCRESAHA.114.302347.
200. Dumitrescu L, Carty CL, Taylor K, Schumacher FR, Hindorf LA, Ambite JL, Anderson G, Best LG, Brown-Gentry K, Bůžková P, Carlson CS, Cochran B, Cole SA, Devereux RB, Duggan D, Eaton CB, Fornage M, Franceschini N, Haessler J, Howard BV, Johnson KC, Laston S, Kolonel LN, Lee ET, MacCluer JW, Manolio TA, Pendergrass SA, Quibera M, Shohet RV, Wilkens LR, Haiman CA, Le Marchand L, Buyske S, Kooperberg C, North KE, Crawford DC. Genetic determinants of lipid traits in diverse populations from the Population Architecture Using Genomics and Epidemiology (PAGE) study. *PLoS Genet*. 2011;7:e1002138. doi: 10.1371/journal.pgen.1002138.
201. Bryant EK, Dressen AS, Bunker CH, Hokanson JE, Hamman RF, Kamboh MI, Demirci FY. A multiethnic replication study of plasma lipoprotein levels-associated SNPs identified in recent GWAS. *PLoS One*. 2013;8:e63469. doi: 10.1371/journal.pone.0063469.
202. Peloso GM, Auer PL, Bis JC, Voorman A, Morrison AC, Stitzel NO, Brody JA, Khetarpal SA, Crosby JR, Fornage M, Isaacs A, Jakobsdottir J, Feitosa MF, Davies G, Huffman JE, Manichaikul A, Davis B, Lohman K, Joon AY, Smith AV, Grove ML, Zanon P, Redon V, Demissie S, Lawson K, Peters U, Carlson C, Jackson RD, Ryckman KK, Mackey RH, Robinson JG, Siscovick DS, Schreiner PJ, Mychaleckyj JC, Pankow JS, Hofman A, Uitterlinden AG, Harris TB, Taylor KD, Stafford JM, Reynolds LM, Marioni RE, Dehghan A, Franco OH, Patel AP, Lu Y, Hindy G, Gottesman O, Bottinger EP, Melander O, Orho-Melander M, Loos RJ, Duga S, Merlini PA, Farrall M, Goel A, Asselta R, Girelli D, Martinelli N, Shah SH, Kraus WE, Li M, Rader DJ, Reilly MP, McPherson R, Watkins H, Ardisson D; NHLBI GO Exome Sequencing Project, Zhang Q, Wang J, Tsai MY, Taylor HA, Correa A, Griswold ME, Lange LA, Starr JM, Rudan I, Eiriksdottir G, Launer LJ, Ordovas JM, Levy D, Chen YD, Reiner AP, Hayward C, Polasek O, Deary IJ, Borecki IB, Liu Y, Gudnason V, Wilson JG, van Duijn CM, Kooperberg C, Rich SS, Psaty BM, Rotter JJ, O'Donnell CJ, Rice K, Boerwinkle E, Kathiresan S, Cupples LA. Association of low-frequency and rare coding-sequence variants with

- blood lipids and coronary heart disease in 56,000 whites and blacks. *Am J Hum Genet.* 2014;94:223–232. doi: 10.1016/j.ajhg.2014.01.009.
203. Fox ER, Musani SK, Barbalic M, Lin H, Yu B, Ogunyankin KO, Smith NL, Kutlar A, Glazer NL, Post WS, Paltoo DN, Dries DL, Farlow DN, Duarte CW, Kardina SL, Meyers KJ, Sun YV, Arnett DK, Patki AA, Sha J, Cui X, Samdarshi TE, Penman AD, Bibbins-Domingo K, Bůžková P, Benjamin EJ, Bluemke DA, Morrison AC, Heiss G, Carr JJ, Tracy RP, Mosley TH, Taylor HA, Psaty BM, Heckbert SR, Cappola TP, Vasan RS. Genome-wide association study of cardiac structure and systolic function in African Americans: the Candidate Gene Association Resource (CARE) study. *Circ Cardiovasc Genet.* 2013;6:37–46. doi: 10.1161/CIRCGENETICS.111.962365.
 204. Akybekova EL, Payne JP, Newton-Cheh C, May WL, Fox ER, Wilson JG, Sarpong DF, Taylor HA, Maher JF. Gene-environment interaction between SCN5A-1103Y and hypokalemia influences QT interval prolongation in African Americans: the Jackson Heart Study. *Am Heart J.* 2014;167:116–122.e1. doi: 10.1016/j.ahj.2013.10.009.
 205. Jones BL, Vyhlidal CA, Bradley-Ewing A, Sherman A, Goggin K. If we would only ask: how Henrietta lacks continues to teach us about perceptions of research and genetic research among African Americans today. *J Racial Ethn Health Disparities.* 2017;4:735–745. doi: 10.1007/s40615-016-0277-1.
 206. Millon Underwood S, Buseh AG, Kelber ST, Stevens PE, Townsend L. Enhancing the participation of African Americans in health-related genetic research: findings of a collaborative academic and community-based research study. *Nurs Res Pract.* 2013;2013:749563. doi: 10.1155/2013/749563.
 207. Ford ES, Capewell S. Proportion of the decline in cardiovascular mortality disease due to prevention versus treatment: public health versus clinical care. *Annu Rev Public Health.* 2011;32:5–22. doi: 10.1146/annurev-publhealth-031210-101211.
 208. Almoudi M, Sun Z. Coronary artery calcium score: re-evaluation of its predictive value for coronary artery disease. *World J Cardiol.* 2012;4:284–287. doi: 10.4330/wjcv.v4.i10.284.
 209. Loria CM, Liu K, Lewis CE, Hulley SB, Sidney S, Schreiner PJ, Williams OD, Bild DE, Detrano R. Early adult risk factor levels and subsequent coronary artery calcification: the CARDIA Study. *J Am Coll Cardiol.* 2007;49:2013–2020. doi: 10.1016/j.jacc.2007.03.009.
 210. Bild DE, Detrano R, Peterson D, Guerci A, Liu K, Shahar E, Ouyang P, Jackson S, Saad MF. Ethnic differences in coronary calcification: the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation.* 2005;111:1313–1320. doi: 10.1161/01.CIR.0000157730.94423.4B.
 211. Manolio TA, Arnold AM, Post W, Bertoni AG, Schreiner PJ, Sacco RL, Saad MF, Detrano RL, Szklo M. Ethnic differences in the relationship of carotid atherosclerosis to coronary calcification: the Multi-Ethnic Study of Atherosclerosis. *Atherosclerosis.* 2008;197:132–138. doi: 10.1016/j.atherosclerosis.2007.02.030.
 212. Wassel CL, Pankow JS, Peralta CA, Choudhry S, Seldin MF, Arnett DK. Genetic ancestry is associated with subclinical cardiovascular disease in African-Americans and Hispanics from the Multi-Ethnic Study of Atherosclerosis. *Circ Cardiovasc Genet.* 2009;2:629–636. doi: 10.1161/CIRCGENETICS.109.876243.
 213. Reiner AP, Ziv E, Lind DL, Nievergelt CM, Schork NJ, Cummings SR, Phong A, Burchard EG, Harris TB, Psaty BM, Kwok PY. Population structure, admixture, and aging-related phenotypes in African American adults: the Cardiovascular Health Study. *Am J Hum Genet.* 2005;76:463–477. doi: 10.1086/428654.
 214. Reiner AP, Carlson CS, Ziv E, Iribarren C, Jaquish CE, Nickerson DA. Genetic ancestry, population sub-structure, and cardiovascular disease-related traits among African-American participants in the CARDIA Study. *Hum Genet.* 2007;121:565–575. doi: 10.1007/s00439-007-0350-2.
 215. Loehr LR, Espeland MA, Sutton-Tyrrell K, Burke GL, Crouse JR 3rd, Herrington DM. Racial differences in endothelial function in post-menopausal women. *Am Heart J.* 2004;148:606–611. doi: 10.1016/j.ahj.2004.04.032.
 216. Heffernan KS, Jae SY, Wilund KR, Woods JA, Fernhall B. Racial differences in central blood pressure and vascular function in young men. *Am J Physiol Heart Circ Physiol.* 2008;295:H2380–H2387. doi: 10.1152/ajpheart.00902.2008.
 217. Ozkor MA, Rahman AM, Murrow JR, Kavtaradze N, Lin J, Manatunga A, Hayek S, Quyyumi AA. Differences in vascular nitric oxide and endothelium-derived hyperpolarizing factor bioavailability in blacks and whites. *Arterioscler Thromb Vasc Biol.* 2014;34:1320–1327. doi: 10.1161/ATVBAHA.113.303136.
 218. D'Agostino RB Sr, Grundy S, Sullivan LM, Wilson P; CHD Risk Prediction Group. Validation of the Framingham coronary heart disease prediction scores: results of a multiple ethnic groups investigation. *JAMA.* 2001;286:180–187.
 219. Lloyd-Jones DM. Cardiovascular risk prediction: basic concepts, current status, and future directions. *Circulation.* 2010;121:1768–1777. doi: 10.1161/CIRCULATIONAHA.109.849166.
 220. McClure LA, Kleindorfer DO, Kissela BM, Cushman M, Soliman EZ, Howard G. Assessing the performance of the Framingham Stroke Risk Score in the Reasons for Geographic and Racial Differences in Stroke cohort. *Stroke.* 2014;45:1716–1720. doi: 10.1161/STROKEAHA.114.004915.
 221. Howard G, Cushman M, Prineas RJ, Howard VJ, Moy CS, Sullivan LM, D'Agostino RB Sr, McClure LA, Pulley L, Safford MM. Advancing the hypothesis that geographic variations in risk factors contribute relatively little to observed geographic variations in heart disease and stroke mortality. *Prev Med.* 2009;49:129–132. doi: 10.1016/j.ypmed.2009.03.004.
 222. Goff DC Jr, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB, Gibbons R, Greenland P, Lackland DT, Levy D, O'Donnell CJ, Robinson J, Schwartz JS, Shero ST, Smith SC, Sorlie P, Stone NJ, Wilson PWF. 2013 Report on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines [data supplement]. *Circulation.* 2014;129(suppl 2):S49–S73. <http://circ.ahajournals.org/lookup/suppl/doi:10.1161/01.cir.0000437741.48606.98/-/DC1>. Accessed August 28, 2017.
 - 222a. Committee on Standards for Developing Trustworthy Clinical Practice Guidelines, Institute of Medicine. *Clinical Practice Guidelines We Can Trust.* Washington, DC: The National Academies Press; 2011.
 223. Kullo IJ, Trejo-Gutierrez JF, Lopez-Jimenez F, Thomas RJ, Allison TG, Mulvagh SL, Arruda-Olson AM, Hayes SN, Pollak AW, Kopecky SL, Hurst RT. A perspective on the new American College of Cardiology/American Heart Association guidelines for cardiovascular risk assessment. *Mayo Clin Proc.* 2014;89:1244–1256. doi: 10.1016/j.mayocp.2014.06.018.
 224. Goff DC Jr, Lloyd-Jones DM. The Pooled Cohort risk equations: black risk matters. *JAMA Cardiol.* 2016;1:12–13. doi: 10.1001/jamacardio.2015.0323.
 225. Centers for Disease Control and Prevention. Interactive atlas of heart disease and stroke tables. 2012. <https://nccd.cdc.gov/dhdspatlas/>. Accessed February 17, 2016.
 226. Muntner P, Colantonio LD, Cushman M, Goff DC Jr, Howard G, Howard VJ, Kissela B, Levitan EB, Lloyd-Jones DM, Safford MM. Validation of the atherosclerotic cardiovascular disease Pooled Cohort risk equations. *JAMA.* 2014;311:1406–1415. doi: 10.1001/jama.2014.2630.
 227. Fox ER, Samdarshi TE, Musani SK, Pencina MJ, Sung JH, Bertoni AG, Xanthakis V, Balfour PC Jr, Shreenivas SS, Covington C, Liebson PR, Sarpong DF, Butler KR, Mosley TH, Rosamond WD, Folsom AR, Herrington DM, Vasan RS, Taylor HA. Development and validation of risk prediction models for cardiovascular events in black adults: the Jackson Heart Study Cohort. *JAMA Cardiol.* 2016;1:15–25. doi: 10.1001/jamacardio.2015.0300.
 228. Taylor HA Jr, Wilson JG, Jones DW, Sarpong DF, Srinivasan A, Garrison RJ, Nelson C, Wyatt SB. Toward resolution of cardiovascular health disparities in African Americans: design and methods of the Jackson Heart Study. *Ethn Dis.* 2005;15(suppl 6):S6–S617.
 229. Howard G, Kissela BM, Kleindorfer DO, McClure LA, Soliman EZ, Judd SE, Rhodes JD, Cushman M, Moy CS, Sands KA, Howard VJ. Differences in the role of black race and stroke risk factors for first vs. recurrent stroke. *Neurology.* 2016;86:637–642.
 230. Sharma A, Colvin-Adams M, Yancy CW. Heart failure in African Americans: disparities can be overcome. *Cleve Clin J Med.* 2014;81:301–311. doi: 10.3949/ccjm.81a.13045.
 231. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA, McBride PE, McMurray JJ, Mitchell JE, Peterson PN, Riegel B, Sam F, Stevenson LW, Tang WH, Tsai EJ, Wilkoff BL. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation.* 2013;128:1810–1852. doi: 10.1161/CIR.0b013e31829e8807.
 232. Exner DV, Dries DL, Domanski MJ, Cohn JN. Lesser response to angiotensin-converting-enzyme inhibitor therapy in black as compared with white patients with left ventricular dysfunction. *N Engl J Med.* 2001;344:1351–1357. doi: 10.1056/NEJM200105033441802.

233. Beta-Blocker Evaluation of Survival Trial Investigators, Eichhorn EJ, Domanski MJ, Krause-Steinrauf H, Bristow MR, Lavori PW. A trial of the beta-blocker bucindolol in patients with advanced chronic heart failure. *N Engl J Med*. 2001;344:1659–1667.
234. Shekelle PG, Rich MW, Morton SC, Atkinson CS, Tu W, Maglione M, Rhodes S, Barrett M, Fonarow GC, Greenberg B, Heidenreich PA, Knabel T, Konstam MA, Steimle A, Warner Stevenson L. Efficacy of angiotensin-converting enzyme inhibitors and beta-blockers in the management of left ventricular systolic dysfunction according to race, gender, and diabetic status: a meta-analysis of major clinical trials. *J Am Coll Cardiol*. 2003;41:1529–1538.
235. Taylor AL, Ziesche S, Yancy C, Carson P, D'Agostino R Jr, Ferdinand K, Taylor M, Adams K, Sabolinski M, Worcel M, Cohn JN; African-American Heart Failure Trial Investigators. Combination of isosorbide dinitrate and hydralazine in blacks with heart failure [published correction appears in *N Engl J Med*. 2005;352:1276]. *N Engl J Med*. 2004;351:2049–2057. doi: 10.1056/NEJMoa042934.
236. McNamara DM, Tam SW, Sabolinski ML, Tobelmann P, Janosko K, Taylor AL, Cohn JN, Feldman AM, Worcel M. Aldosterone synthase promoter polymorphism predicts outcome in African Americans with heart failure: results from the A-HeFT Trial. *J Am Coll Cardiol*. 2006;48:1277–1282. doi: 10.1016/j.jacc.2006.07.030.
237. Lewis LM, Ogedegbe C, Ogedegbe G. Enhancing adherence of antihypertensive regimens in hypertensive African-Americans: current and future prospects. *Expert Rev Cardiovasc Ther*. 2012;10:1375–1380. doi: 10.1586/erc.12.138.
238. Egan BM, Laken MA. Is blood pressure control to less than 140/less than 90 mmHg in 50% of all hypertensive patients as good as we can do in the USA: or is this as good as it gets? *Curr Opin Cardiol*. 2011;26:300–307. doi: 10.1097/HCO.0b013e3283474c20.
239. Harman J, Walker ER, Charbonneau V, Akybekova EL, Nelson C, Wyatt SB. Treatment of hypertension among African Americans: the Jackson Heart Study. *J Clin Hypertens (Greenwich)*. 2013;15:367–374. doi: 10.1111/jch.12088.
240. Johnson JA. Ethnic differences in cardiovascular drug response: potential contribution of pharmacogenetics. *Circulation*. 2008;118:1383–1393. doi: 10.1161/CIRCULATIONAHA.107.704023.
241. Fuchs FD. Why do black Americans have higher prevalence of hypertension? An enigma still unsolved. *Hypertension*. 2011;57:379–380. doi: 10.1161/HYPERTENSIONAHA.110.163196.
242. Ramsay LE, Williams B, Johnston GD, MacGregor GA, Poston L, Potter JF, Poulter NR, Russell G. British Hypertension Society guidelines for hypertension management 1999: summary. *BMJ*. 1999;319:630–635.
243. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, Lackland DT, LeFevre ML, MacKenzie TD, Ogedegbe O, Smith SC Jr, Svetkey LP, Taler SJ, Townsend RR, Wright JT Jr, Narva AS, Ortiz E. 2014 Evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8) [published correction appears in *JAMA*. 2014;311:1809]. *JAMA*. 2014;311:507–520. doi: 10.1001/jama.2013.284427.
244. Wright JT Jr, Fine LJ, Lackland DT, Ogedegbe G, Dennison-Himmelfarb CR. Evidence supporting a systolic blood pressure goal of less than 150 mm Hg in patients aged 60 years or older: the minority view. *Ann Intern Med*. 2014;160:499–503. doi: 10.7326/M13-2981.
245. Laakkanen JA, Kurl S, Salonen JT. Cardiorespiratory fitness and physical activity as risk predictors of future atherosclerotic cardiovascular diseases. *Curr Atheroscler Rep*. 2002;4:468–476.
246. Brewster LM, van Montfrans GA, Kleijnen J. Systematic review: antihypertensive drug therapy in black patients. *Ann Intern Med*. 2004;141:614–627.
247. Preston RA, Materson BJ, Reda DJ, Williams DW, Hamburger RJ, Cushman WC, Anderson RJ. Age-race subgroup compared with renin profile as predictors of blood pressure response to antihypertensive therapy: Department of Veterans Affairs Cooperative Study Group on Antihypertensive Agents. *JAMA*. 1998;280:1168–1172.
248. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr*. 1982;36:936–942.
249. Materson BJ, Reda DJ, Cushman WC, Massie BM, Freis ED, Kochar MS, Hamburger RJ, Fye C, Lakshman R, Gottdiener J, Ramirez EJ, Henderson WG; Department of Veterans Affairs Cooperative Study Group on Antihypertensive Agents. Single-drug therapy for hypertension in men: a comparison of six antihypertensive agents with placebo: the Department of Veterans Affairs Cooperative Study Group on Antihypertensive Agents [published correction appears in *N Engl J Med*. 1994;330:1689]. *N Engl J Med*. 1993;328:914–921.
250. Agodoa LY, Appel L, Bakris GL, Beck G, Bourgoignie J, Briggs JP, Charleston J, Cheek D, Cleveland W, Douglas JG, Douglas M, Dowie D, Faulkner M, Gabriel A, Gassman J, Greene T, Hall Y, Hebert L, Hiremath L, Jamerson K, Johnson CJ, Kopple J, Kusek J, Lash J, Lea J, Lewis JB, Lipkowitz M, Massry S, Middleton J, Miller ER 3rd, Norris K, O'Connor D, Ojo A, Phillips RA, Pogue V, Rahman M, Randall OS, Rostand S, Schulman G, Smith W, Thornley-Brown D, Tisher CC, Toto RD, Wright JT Jr, Xu S; African American Study of Kidney Disease and Hypertension (AASK) Study Group. Effect of ramipril vs amlodipine on renal outcomes in hypertensive nephrosclerosis: a randomized controlled trial. *JAMA*. 2001;285:2719–2728.
251. Jaffe MG, Lee GA, Young JD, Signey S, Go AS. Improved blood pressure control associated with a large-scale hypertension program. *JAMA*. 2013;310:699–705.
252. Go AS, Bauman MA, Coleman King SM, Fonarow GC, Lawrence W, Williams KA, Sanchez E. An effective approach to high blood pressure control: a science advisory from the American Heart Association, the American College of Cardiology, and the Centers for Disease Control and Prevention [published correction appears in *Hypertension*. 2014;63:e175]. *Hypertension*. 2014;63:878–885. doi: 10.1161/HYP.0000000000000003.
253. Kotchen TA, Cowley AW Jr, Liang M. Ushering hypertension into a new era of precision medicine. *JAMA*. 2016;315:343–344.
254. Williams DR. Miles to go before we sleep: racial inequities in health. *J Health Soc Behav*. 2012;53:279–295. doi: 10.1177/0022146512455804.
255. Harper S, Lynch J, Smith GD. Social determinants and the decline of cardiovascular diseases: understanding the links. *Annu Rev Public Health*. 2011;32:39–69. doi: 10.1146/annurev-publhealth-031210-101234.
256. Kreatsoulas C, Anand SS. The impact of social determinants on cardiovascular disease. *Can J Cardiol*. 2010;26(suppl C):8C–13C.
257. Everson-Rose SA, Lewis TT. Psychosocial factors and cardiovascular diseases. *Annu Rev Public Health*. 2005;26:469–500. doi: 10.1146/annurev.publhealth.26.021304.144542.
258. Havranek EP, Mujahid MS, Barr DA, Blair IV, Cohen MS, Cruz-Flores S, Davey-Smith G, Dennison-Himmelfarb CR, Lauer MS, Lockwood DW, Rosal M, Yancy CW; on behalf of the American Heart Association Council on Quality of Care and Outcomes Research, Council on Epidemiology and Prevention, Council on Cardiovascular and Stroke Nursing, Council on Lifestyle and Cardiometabolic Health, and Stroke Council. Social determinants of risk and outcomes for cardiovascular disease: a scientific statement from the American Heart Association. *Circulation*. 2015;132:873–898. doi: 10.1161/CIR.0000000000000228.
259. DeNavas-Walt C, Proctor BD. *U.S. Census Bureau, Current Population Reports, P60-252, Income and Poverty in the United States: 2014*. Washington, DC: US Government Printing Office; 2015.
260. Baker EA, Schootman M, Barnidge E, Kelly C. The role of race and poverty in access to foods that enable individuals to adhere to dietary guidelines. *Prev Chronic Dis*. 2006;3:A76.
261. Yancey AK, Cole BL, Brown R, Williams JD, Hillier A, Kline RS, Ashe M, Grier SA, Backman D, McCarthy WJ. A cross-sectional prevalence study of ethnically targeted and general audience outdoor obesity-related advertising. *Milbank Q*. 2009;87:155–184. doi: 10.1111/j.1468-0009.2009.00551.x.
262. Dias JJ, Whitaker RC. Black mothers' perceptions about urban neighborhood safety and outdoor play for their preadolescent daughters. *J Health Care Poor Underserved*. 2013;24:206–219. doi: 10.1353/hpu.2013.0018.
263. Frierson GM, Howard EN, DeFina LE, Powell-Wiley TM, Willis BL. Effect of race and socioeconomic status on cardiovascular risk factor burden: the Cooper Center Longitudinal Study. *Ethn Dis*. 2013;23:35–42.
264. Winkleby MA, Kraemer HC, Ahn DK, Varady AN. Ethnic and socioeconomic differences in cardiovascular disease risk factors: findings for women from the Third National Health and Nutrition Examination Survey, 1988–1994. *JAMA*. 1998;280:356–362.
265. Kershaw KN, Albrecht SS. Racial/ethnic residential segregation and cardiovascular disease risk. *Curr Cardiovasc Risk Rep*. 2015;9:10.
266. Kershaw KN, Osypuk TL, Do DP, De Chavez PJ, Diez Roux AV. Neighborhood-level racial/ethnic residential segregation and incident cardiovascular disease: the Multi-Ethnic Study of Atherosclerosis. *Circulation*. 2015;131:141–148. doi: 10.1161/CIRCULATIONAHA.114.011345.

267. Casper M, Nwaise I, Croft JB, Hong Y, Fang J, Greer S. Geographic disparities in heart failure hospitalization rates among Medicare beneficiaries. *J Am Coll Cardiol*. 2010;55:294–299. doi: 10.1016/j.jacc.2009.10.021.
268. Bonora E, Kiechl S, Willeit J, Oberhollenzer F, Egger G, Bonadonna RC, Muggeo M; Bruneck Study. Carotid atherosclerosis and coronary heart disease in the metabolic syndrome: prospective data from the Bruneck study. *Diabetes Care*. 2003;26:1251–1257.
269. Howard G. Ancel Keys Lecture: adventures (and misadventures) in understanding (and reducing) disparities in stroke mortality. *Stroke*. 2013;44:3254–3259. doi: 10.1161/STROKEAHA.113.002113.
270. Lewis TT, Williams DR, Tamene M, Clark CR. Self-reported experiences of discrimination and cardiovascular disease. *Curr Cardiovasc Risk Rep*. 2014;8:365. doi: 10.1007/s12170-013-0365-2.
271. Sampson UK, Kaplan RM, Cooper RS, Diez Roux AV, Marks JS, Engelgau MM, Peprah E, Mishoe H, Boulware LE, Felix KL, Califf RM, Flack JM, Cooper LA, Gracia JN, Henderson JA, Davidson KW, Krishnan JA, Lewis TT, Sanchez E, Luban NL, Vaccarino V, Wong WF, Wright JT Jr, Meyers D, Ogedegbe OG, Presley-Cantrell L, Chambers DA, Belis D, Bennett GC, Boyington JE, Creazzo TL, de Jesus JM, Krishnamurti C, Lowden MR, Punturieri A, Shero ST, Young NS, Zou S, Mensah GA. Reducing health inequities in the U.S.: recommendations from the NHLBI's Health Inequities Think Tank Meeting. *J Am Coll Cardiol*. 2016;68:517–524. doi: 10.1016/j.jacc.2016.04.059.
272. Black LL, Johnson R, VanHoose L. The relationship between perceived racism/discrimination and health among black American women: a review of the literature from 2003 to 2013. *J Racial Ethn Health Disparities*. 2015;2:11–20. doi: 10.1007/s40615-014-0043-1.
273. Lewis TT, Cogburn CD, Williams DR. Self-reported experiences of discrimination and health: scientific advances, ongoing controversies, and emerging issues. *Annu Rev Clin Psychol*. 2015;11:407–440. doi: 10.1146/annurev-clinpsy-032814-112728.
274. Dolezsar CM, McGrath JJ, Herzog AJ, Miller SB. Perceived racial discrimination and hypertension: a comprehensive systematic review. *Health Psychol*. 2014;33:20–34. doi: 10.1037/a0033718.
275. Stepanikova I, Baker EH, Simoni ZR, Zhu A, Rutland SB, Sims M, Wilkinson LL. The role of perceived discrimination in obesity among African Americans. *Am J Prev Med*. 2017;52:S77–S85. doi: 10.1016/j.amepre.2016.07.034.
276. Kershaw KN, Lewis TT, Diez Roux AV, Jenny NS, Liu K, Penedo FJ, Carnethon MR. Self-reported experiences of discrimination and inflammation among men and women: the Multi-Ethnic Study of Atherosclerosis. *Health Psychol*. 2016;35:343–350. doi: 10.1037/hea0000331.
277. Vadiello M, Mattei J. Perceived weight discrimination and 10-year risk of allostatic load among US adults [published correction appears in *Ann Behav Med*. 2017;51:105]. *Ann Behav Med*. 2017;51:94–104. doi: 10.1007/s12160-016-9831-7.
278. Everson-Rose SA, Lutsey PL, Roetker NS, Lewis TT, Kershaw KN, Alonso A, Diez Roux AV. Perceived discrimination and incident cardiovascular events: the Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol*. 2015;182:225–234. doi: 10.1093/aje/kwv035.
279. Dubois CM, Beach SR, Kashdan TB, Nyer MB, Park ER, Celano CM, Huffman JC. Positive psychological attributes and cardiac outcomes: associations, mechanisms, and interventions. *Psychosomatics*. 2012;53:303–318. doi: 10.1016/j.psych.2012.04.004.
280. Klainin-Yobas P, Ng SH, Stephen PD, Lau Y. Efficacy of psychosocial interventions on psychological outcomes among people with cardiovascular diseases: a systematic review and meta-analysis. *Patient Educ Couns*. 2016;99:512–521. doi: 10.1016/j.pec.2015.10.020.
281. Macintosh T, Desai MM, Lewis TT, Jones BA, Nunez-Smith M. Socially-assigned race, healthcare discrimination and preventive healthcare services. *PLoS One*. 2013;8:e64522. doi: 10.1371/journal.pone.0064522.
282. Bennett GG, Wolin KY, Goodman M, Samplin-Salgado M, Carter P, Dutton S, Hill R, Emmons K. Attitudes regarding overweight, exercise, and health among blacks (United States). *Cancer Causes Control*. 2006;17:95–101. doi: 10.1007/s10552-005-0412-5.
283. Ard JD, Fitzpatrick S, Desmond RA, Sutton BS, Pisu M, Allison DB, Franklin F, Baskin ML. The impact of cost on the availability of fruits and vegetables in the homes of schoolchildren in Birmingham, Alabama. *Am J Public Health*. 2007;97:367–372. doi: 10.2105/AJPH.2005.080655.
284. Kumanyika SK, Wadden TA, Shults J, Fassbender JE, Brown SD, Bowman MA, Brake V, West W, Frazier J, Whitt-Glover MC, Kallan MJ, Desnouvee E, Wu X. Trial of family and friend support for weight loss in African American adults. *Arch Intern Med*. 2009;169:1795–1804. doi: 10.1001/archinternmed.2009.337.
285. Samuel-Hodge CD, Gizlice Z, Cai J, Brantley PJ, Ard JD, Svetkey LP. Family functioning and weight loss in a sample of African Americans and whites. *Ann Behav Med*. 2010;40:294–301. doi: 10.1007/s12160-010-9219-z.
286. Gordon-Larsen P. Obesity-related knowledge, attitudes, and behaviors in obese and non-obese urban Philadelphia female adolescents. *Obes Res*. 2001;9:112–118. doi: 10.1038/oby.2001.14.
287. Kumanyika SK, Morssink C, Agurs T. Models for dietary and weight change in African-American women: identifying cultural components. *Ethn Dis*. 1992;2:166–175.
288. Kumanyika S, Wilson JF, Guilford-Davenport M. Weight-related attitudes and behaviors of black women. *J Am Diet Assoc*. 1993;93:416–422.
289. Fitzgibbon ML, Tussing-Humphreys LM, Porter JS, Martin IK, Odoms-Young A, Sharp LK. Weight loss and African-American women: a systematic review of the behavioural weight loss intervention literature. *Obes Rev*. 2012;13:193–213. doi: 10.1111/j.1467-789X.2011.00945.x.
290. Kong A, Tussing-Humphreys LM, Odoms-Young AM, Stolley MR, Fitzgibbon ML. Systematic review of behavioural interventions with culturally adapted strategies to improve diet and weight outcomes in African American women. *Obes Rev*. 2014;15(suppl 4):62–92. doi: 10.1111/obr.12203.
291. Lancaster KJ, Carter-Edwards L, Grilo S, Shen C, Schoenthaler AM. Obesity interventions in African American faith-based organizations: a systematic review. *Obes Rev*. 2014;15(suppl 4):159–176. doi: 10.1111/obr.12207.